

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>C07D 257/02, 401/12, 409/12, 403/12, 471/06, 401/06, 473/00, A61K 31/395 //</b> <b>(C07D 471/06, 257:00, 221:00)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/02870</b> <b>(43) International Publication Date:</b> 20 January 2000 (20.01.00)
<b>(21) International Application Number:</b> PCT/CA99/00619 <b>(22) International Filing Date:</b> 8 July 1999 (08.07.99)  <b>(30) Priority Data:</b> 09/111,895 8 July 1998 (08.07.98) US  <b>(71) Applicant (for all designated States except US):</b> ANORMED INC. [CA/CA]; No. 200, 20353 - 64th Avenue, Surrey, British Columbia V2Y 1N5 (CA).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BRIDGER, Gary, James [GB/US]; 445 S. Garden Street, Bellingham, WA 98225 (US). BOEHRINGER, Eva, Maria [CA/CA]; 404 - 1558 Fir Street, White Rock, British Columbia V4B 4B7 (CA). WANG, Zhongren [CA/CA]; #218, 5380 - 208 Street, Langley, British Columbia V3A 2C2 (CA). SCHOLS, Dominique [BE/BE]; Kiersveld 9, B-3020 Herent (BE). SKERLJ, Renato, Tony [NZ/US]; 9062 Shearwater Road, Blaine, WA 98230 (US). BOGUCKI, David, Earl [CA/CA]; 203, 16071 - 82nd Avenue, Surrey, British Columbia V3S 2L6 (CA).		<b>(74) Agents:</b> ROBINSON, J., Chistopher et al.; Fetherstonhaugh & Co., Suite 2200, 650 West Georgia Street, Box 11560, Vancouver, British Columbia V6B 4N8 (CA).  <b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ANTIVIRAL MACROCYCLIC COMPOUNDS  <b>(57) Abstract</b>  The present invention is drawn to novel antiviral compounds, pharmaceutical compositions and their use. More specifically this invention is drawn to derivatives of monocyclic polyamines which have activity in standard tests against HIV- or FIV-infected cells as well as other biological activity related to binding of ligands to chemokine receptors that mediate a number of mammalian embryonic developmental processes.		

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## ANTIVIRAL MACROCYCLIC COMPOUNDS

The present invention is drawn to novel antiviral compounds, pharmaceutical compositions and their use. More specifically this invention is drawn to derivatives of monocyclic polyamines which have activity in standard tests against HIV-infected cells as well as other biological activity related to binding of ligands to chemokine receptors that mediate a number of mammalian embryonic developmental processes. The present invention further includes methods of treating various diseases mediated by chemokine receptor-ligand binding.

### BACKGROUND OF THE INVENTION

Approximately 40 human chemokines have been described, that function, at least in part, by modulating a complex and overlapping set of biological activities important for the movement of lymphoid cells and extravasation and tissue infiltration of leukocytes in response to inciting agents (See, for example: P. Ponath, *Exp. Opin. Invest. Drugs*, 7:1-18, 1998). These chemotactic cytokines, or chemokines, constitute a family of proteins, approximately 8-10 kDa in size. Chemokines appear to share a common structural motif, that consists of 4 conserved cysteines involved in maintaining tertiary structure. There are two major subfamilies of chemokines: the "CC" or  $\beta$ -chemokines and the "CXC" or  $\alpha$ -chemokines. The receptors of these chemokines are classified based upon the chemokine that constitutes the receptor's natural ligand. Receptors of the  $\beta$ -chemokines are designated "CCR"; while those of the  $\alpha$ -chemokines are designated "CXCR".

Chemokines are considered to be principal mediators in the initiation and maintenance of inflammation. More specifically, chemokines have been found to play an important role in the regulation of endothelial cell function, including proliferation, migration and differentiation during angiogenesis and re-endothelialization after injury (Gupta *et al.*, *J. Biolog. Chem.*, 7:4282-4287, 1998). Two specific chemokines have been implicated in the etiology of infection by human immunodeficiency virus (HIV).

In most instances, HIV initially binds via its gp120 envelope protein to the CD4 receptor of the target cell. A conformational change appears to take place in the gp120 which results in

its subsequent binding to a chemokine receptor, such as CCR-5 (Wyatt *et al.*, *Science*, 280:1884-1888 (1998)). HIV-1 isolates arising subsequently in the infection bind to the CXCR-4 chemokine receptor. In view of the fact that the feline immunodeficiency virus, another related retrovirus, binds to a chemokine receptor without needing to bind first to the CD4 receptor, suggests that chemokine receptors may be the primordial obligate receptors for immunodeficiency retroviruses (Richardson *et al.*, *J. Virol.* 73:661 (1999)).

Following the initial binding by HIV to CD4, virus-cell fusion results, which is mediated by members of the chemokine receptor family, with different members serving as fusion cofactors for macrophage-tropic (M-tropic) and T cell line-tropic (T-tropic) isolates of HIV-1 (Carroll *et al.*, *Science*, 276: 273-276 1997). During the course of infection within a patient, it appears that a majority of HIV particles shift from the M-tropic to the more aggressive T-tropic viral phenotype (Miedema *et al.*, *Immune. Rev.*, 140:35 (1994)). Curiously, the M-tropic viral phenotype correlates with the virus's ability to enter the cell following binding of the CCR-5 receptor, while the T-tropic viral phenotype correlates with viral entry into the cell following binding and membrane fusion with the CXCR-4 receptor. Clinically observations suggest that patients who possess genetic mutations in the CCR-5 appear resistant or less susceptible to HIV infection.

However, the binding of chemokine receptors to their natural ligands appears to serve a more evolutionary and central role than only as mediators of HIV infection. The chemokine receptor, CXCR-4 has been found to be essential for the vascularization of the gastrointestinal tract (Tachibana *et al.*, *Nature*, 393:591-594 (1998)) as well as haematopoiesis and cerebellar development (Zou *et al.*, *Nature*, 393:591-594 (1998)). Interference with any of these important functions served by the binding of pre-B-cell growth-stimulating factor/stromal derived factor (PBSF/SDF-1) to the CXCR-4 chemokine receptor results in lethal deficiencies in vascular development, haematopoiesis and cardiogenesis. Similarly, fetal cerebellar development appears to rely upon the effective functioning of CXCR-4 in neuronal cell migration and patterning in the central nervous system. This G-protein-coupled chemokine receptor appears to play a critical role in ensuring the necessary patterns of migration of granule cells in the cerebellar anlage. Interactions of SDF-1 and CXCR4 are also important in maintaining B-cell lineage and in retaining stem cells in bone marrow (Peled *et al.*, *Science* 283: 845 (1999); Springer *et al.*, *Immunity* 10:463 (1999))

In attempting to better understand the relationship between chemokines and their receptors, recent experiments to block the binding of HIV to the CXCR-4 chemokine receptor were carried out through the use of monoclonal antibodies or small molecules that appear to suggest a useful therapeutic strategy (Schols *et al.*, *J. Exp. Med.* 186:1383-1388 (1997); Schols *et al.*, *Antiviral Research* 35:147-156 (1997)). Small molecules, such as bicyclams, appear to specifically interfere with the CXCR-4 binding and not CCR-5 binding (Donzella *et al.*, *Nature Medicine*, 4:72-77 (1998)). These experiments demonstrated interference with HIV entry and membrane fusion into the target cell *in vitro*. Additional experiments monitoring the calcium flux or  $Ca^{2+}$  mobilization assay demonstrated that a bicyclam also functioned as an antagonist to signal transduction resulting from the binding of stromal derived factor or SDF-1 $\alpha$ , the natural chemokine to CXCR-4.

Further, the etiology or association of chemokine receptor binding in the proliferation of glioblastoma tumor cells has been reported by Sehgal *et al.*, *J. of Surg. Oncolo.* 69:99-104 (1998) ("Sehgal I") and Sehgal *et al.*, *J. of Surg. Oncolo.* 69:239-248 (1998) ("Sehgal II"). The role of CXCR4 of its binding to its receptor appears to play a significant role in the formation and/or proliferation of glioblastoma cells. The inhibition of the binding by CXCR4 to its natural receptor ligand by compounds of the present invention, such as AMD 3100, offer a new drug in the treatment tumors of central nervous system that are mediated or associated with chemokines, such as CXCR4.

Additionally, CXC chemokines have been found to regulate or are associated with the regulation of angiogenesis in non-small cell lung cancer (see: Arenberg, *et al.*, *J. of Leukocyte Biol.*; 62:554-562 (1997); and Moore *et al.* *TCM*, vol 8(2): 51-58 (1998) Elsevier Science, Inc.). The role of CXC chemokines and the binding to their respective receptors appear to play a significant role in the formation and/or proliferation of non-small cell lung cancer. The inhibition of the binding of these CXC chemokines to their natural receptor ligands by compounds of the present invention, such as AMD 3100, offer a new drug in the treatment tumors such as non-small cell lung cancer that are mediated or associated with increased levels of chemokines.


U.S. Pat. No. 5,583,131, U.S. Pat. No. 5,698,546 and U.S. Pat. No. 5,817,807 disclose cyclic compounds that are active against HIV-1 and HIV-2 in *in vitro* tests. We have now discovered that these compounds exhibit anti-HIV (anti-human immunodeficiency virus) and

anti-FIV (anti-feline immunodeficiency virus) activities due to their binding to the chemokine receptor 4 (CXCR-4 or Fusin receptor), expressed on the surface of certain cells of the immune system (Este *et al.*, *Mol. Pharmacol.* 55:67 (1999); Egberink *et al.*, *J. Virol in press* (1999)). This competitive binding thereby protects these target cells from infection by HIV which utilize the CXCR-4 receptor for entry. We have discovered that the disclosed compounds also antagonize the binding, signaling and chemotactic effects of the natural CXC-chemokine for CXCR-4, stromal cell-derived factor 1 $\alpha$  (SDF-1 $\alpha$ ). Herein, we further disclose novel compounds that demonstrate protective effects against HIV infection of target cells by inhibition of binding *in vitro* to the CC-5 receptor (CCR-5).

#### SUMMARY OF THE INVENTION

The present invention provides novel compounds, that demonstrate protective effects on target cells from HIV infection as well as demonstrate other biological activities related to the ability of these compounds to inhibit the binding by the natural ligand to its chemokine receptor.

Accordingly, the present invention provides a macrocyclic compound of formula I:



wherein V is a cyclic polyamine moiety having a total of 9 to 24 members and from 2 to 6, but preferably from 3 to 6, optionally substituted amino nitrogens optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic ring;

R<sup>1</sup> to R<sup>7</sup> may be the same or different and are independently selected from hydrogen or straight, branched or cyclic C<sub>1-6</sub> alkyl;

R<sup>8</sup> is a heterocyclic group, a substituted aromatic group, or a mercaptan group;

Ar is an aromatic or heteroaromatic ring each optionally substituted at single or multiple positions with electrons-donating or withdrawing groups;

x is 1 or 2;

and the acid addition salts and metal complexes thereof.

Preferably V is a 14- to 17 membered fused or unfused ring system, such as a cyclam

system or a 4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene system or a derivative thereof, and especially a cyclam system or derivative thereof. The moiety V may be substituted at C or N non-linking atoms, suitably by hydroxyl, alkoxy, thiol, thioalkyl or any other atom or group which does not adversely affect the activity or toxicity of the compounds but may reduce the basicity of the amines, for example halogen, nitro, carboxy, carboxamido, sulphonic acid or phosphate. Suitably the fused aromatic or heteroaromatic ring is phenyl, pyridine, pyrimidine, pyrazine, imidazole or thiazole. Preferably, the fused aromatic or heteroaromatic ring is phenyl or pyridine.

10 Preferably R<sup>1</sup> to R<sup>7</sup> are each hydrogen.

Preferably R<sup>8</sup> is selected from pyridine, pyrimidine, pyrazine, imidazole, thiophene, thiophenyl, aminobenzyl, piperidinyl, piperazinyl or a mercaptan group.

15 Preferably Ar is phenyl. Preferred substituents are alkyl, aryl, amino, alkoxy, hydroxy, halogen, carboxyl and carboxamido.


The invention also includes what may be termed as "pro-drug", that is protected forms of the compounds, which release the compound after administration to a patient. For example, the compound may carry a protective groups which is split off by hydrolysis in body fluids e.g. in the bloodstream, thus releasing active compound or are oxidized or reduced in body fluids to release the compound. A discussion on pro-drugs may be found in "Smith and Williams' Introduction to the Principles of Drug Design", H.J. Smith, Wright, Second Edition, London 1988.

25 Acid addition salts, for example hydrochlorides, and non-toxic labile metal complexes of compounds of formula I are also active compounds according to the present invention. Non-toxic in the present tense has to be considered with reference to the prognosis for the infected patient without treatment. Copper and zinc complexes are preferred although other metals such as nickel may be considered, whereas less labile metals such as cobalt and rhodium are less preferred because of likely lower selectivity.

30 Compounds of formula I are novel. Accordingly, a further aspect of the invention

provides a process for the preparation of a compound of formula I which comprises the following steps:

- (i) nucleophilic attack by the cyclic polyamine V having a single unprotected amine nitrogen, all other amine nitrogen atoms being protected, on an excess of a compound of formula II




- wherein  $R^1$  to  $R^4$  and Ar are as hereinbefore defined, and each Y is an active substituent which can be displaced by the unprotected nitrogen of polyamine V and is preferably selected from Br, Cl, I, methane sulphonate, 4-toluenesulphonate, trifluoromethane sulphonate.

- It is well within the capabilities and knowledge of the skilled synthetic chemist to protect the amine nitrogens of cyclic polyamines, and it is preferred to use substitution by methanesulphonyl and/or toluenesulphonyl and/or diethoxyphosphoryl (see: Bridger *et al*, *J. Med. Chem.*, 38:366-378 (1995); Bridger *et al*, U.S. Pat. 5,583,131 or Bridger *et al*, U.S. Pat. 5,698,546) and/or nitrobenzenesulfonyl (Fukuyama *et al.*, *Tetrahedron Letters* 1995, 36, 6373-6374).

- The protected polyamine V is firstly reacted with a 5-to 10-fold excess of a compound of formula II in a solvent such as acetonitrile or dimethylformamide, tetrahydrofuran or dioxane and in the presence of a base, for example sodium carbonate or potassium carbonate. The reaction generally proceeds at room temperature to elevated temperature to give a cyclic polyamine in which all amine nitrogens are protected. In general, a mixture of products will be obtained and we have found that the product can conveniently be purified by silica gel chromatography or crystallization.

Nucleophilic attack of a compound of formula III



wherein  $R^5$  to  $R^8$  and x are as hereinbefore defined on the product of the reaction



described at (I) above, and subsequently de-protecting the amine nitrogens. The reaction with an excess of a compound of formula III is carried out under similar conditions to the reaction with the polyamine V.

The de-protection step is suitably carried out by re-fluxing the protected molecule in a mixture of aqueous HBr and acetic acid or concentrated sulphuric acid, or in the case of diethoxyphosphoryl in the presence of gaseous hydrogen chloride or gaseous hydrogen bromide in acetic acid; in the case of nitrobenzenesulfonyl deprotection, a mercaptan such as thiophenol or mercaptoacetic acid in the presence of a suitable base such as potassium carbonate, cesium carbonate, sodium hydroxide or lithium hydroxide in a solvent such as dimethylformamide, acetonitrile, tetrahydrofuran or dioxane is used. This reaction generally proceeds at room temperature to elevated temperatures to give a polyamine in which the nitrogens are deprotected. Alternatively, and accordingly, a further aspect of the invention provides a process for the preparation of compounds of Formula I which comprises the following steps:

(i) nucleophilic attack by the cyclic polyamine V having a single unprotected amine nitrogen, all other amine nitrogens being protected, with an excess of a compound of formula (IV)




Wherein  $R^1$  to  $R^4$  and  $x$ ,  $R^6$  to  $R^8$  and  $Ar$  are as hereinbefore defined and  $Y$  is an active substituent which can be displaced by the unprotected nitrogen of cyclic polyamine V as hereinbefore defined. In this case, the intended substituent  $R^5$  is hydrogen but for convenience, the nitrogen is protected as a nitrobenzenesulfonyl or diethoxyphosphoryl group.

The protected polyamine V is first reacted with a compound of formula IV using similar conditions to the reactions with compounds of formula II and formula III as described above and the product of this reaction is subjected to deprotection of the amine nitrogens on the polyamine and at  $R^5$ .

The deprotection steps were carried out as described above. For convenience, a sequential combination of these deprotection reactions may be used when a mixture of any of: methanesulphonyl; toluenesulphonyl; diethoxyphosphoryl; or nitrobenzenesulfonyl groups are present.

The novel compounds further comprise a macrocyclic compound of general formula V:



where  $V^2$  is a cyclic polyamine moiety having a total of 9 to 24 members and from 2 to 6, but preferably from 3 to 6, optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic ring; where  $R^9$  and  $R^{10}$  may be the same or different and are independently selected from hydrogen or straight, branched or cyclic  $C_{1-6}$  alkyl; further, where  $Ar^2$  is an aromatic, fused aromatic, heterocyclic or fused heterocyclic ring each optionally substituted at single or multiple positions with electron-donating or withdrawing groups and/or aromatic and heterocyclic groups and their alkyl derivatives thereof; and the acid addition salts and metal complexes.

These novel compounds have demonstrated anti-HIV activity in an *in vitro* screen assay as presented in Table 1. These novel compounds have also demonstrated biological activity in inhibiting CXCR-4 specific monoclonal antibody (12G5) from binding to CXCR-4 on SUP-T1 cells by AMD compounds. These data are shown in Table 2 for AMD3100 (1,1'-[1,4-phenylenebis(methylene)]bis-1,4,8,11-tetraazacyclotetradecane), AMD3465 (N-[1,4,8,11-tetraazacyclotetra-decanyl-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine) and six new compounds: AMD 7049; AMD 7050; AMD 7051; AMD 7058; AMD 7059; and AMD 7063.

Data showing inhibition of the increased  $Ca^{2+}$  flux induced by SDF-1 in SUP-T1 cells (inhibition of signal transduction) by AMD compounds are shown in Table 3 for AMD3100, AMD3465 and compounds: AMD 7049; AMD 7050; AMD 7051; AMD 7058; AMD 7059; and AMD 7063.

Several novel compounds also inhibited infection of the cell line U87.CD4.CCR5 by the M-tropic HIV-1 strain BaL, which exclusively utilizes the CCR-5 co-receptor for entry. These data are shown in Table 4.

The experimental procedures for the mAb binding assay, the inhibition of  $Ca^{2+}$  flux, and inhibition of infection by the HIV-1 BaL strain in U87.CD4.CCR5 cells would be readily understood by the skilled artisan. For example, see: Schols *et al.*, *J. Exp. Med.* 186:1383-1388 (1997); Schols *et al.*, *Antiviral Research* 35:147-156 (1997); and Donzella *et al.*, *Nature*

*Medicine*, 4:72-77 (1998). Also, the characterization of the CXCR-4 specific monoclonal antibody 12G5 is taught by Hoxie *et al.*, *Cell*, 87:745-756 (1996).

Citation of the above documents is not intended as an admission that any of the foregoing is pertinent prior art. All statements as to the date or representation as to the contents  
5 of these documents is based on the information available to the applicants and does not constitute any admission as to the correctness of the dates or contents of these documents. Further, all documents referred to throughout this application are incorporated in their entirety by reference herein.

Having now generally described the invention, the same will be more readily understood  
10 through reference to the following examples which are provided by way of illustration, and are not intended to be limiting of the present invention, unless specified.

As mentioned above, the compounds of the invention have activity against viral infections, especially retrovirus infections and specifically HIV. Accordingly, a further aspect of the invention provides a compound of formula I or formula V for use in medicine. More  
15 specifically, there is provided the use of a compound of formula I or formula V in the manufacture of a medicament for the treatment of HIV-infected patients. In the alternative, there is provided a method of treating an HIV-infected patient comprising administering to said patient, a pharmaceutically effective amount of a compound of formula I or formula V. Although compounds of formula I or formula V could be administered as the raw material, it is  
20 preferable to present them in the form of a pharmaceutical composition comprising a compound of formula I or formula V as active ingredient in admixture with a pharmaceutically acceptable diluent or carrier and optionally one or more other therapeutic ingredients, such compositions providing a further aspect of the invention.

In all aspects of the invention, it is understood that meso forms, enantiomers and  
25 resolved optically active forms of the compounds of formula I or formula V are also included. Also, it is to be considered within the invention, compounds of formula I or formula V diluted with non-toxic or other active substances.

#### BRIEF DESCRIPTION OF THE DRAWINGS

30 Fig. 1, shows the structural formula of compound AMD 3465.

Fig. 2, shows the structural formula of compound AMD 3538.

- Fig. 3; shows the structural formula of compound AMD 3500.
- Fig. 4, shows the structural formula of compound AMD 3499.
- Fig. 5; shows the structural formula of compound AMD 3498.
- Fig. 6; shows the structural formula of compound AMD 3497.
- 5 Fig. 7; shows the structural formula of compound AMD 3516.
- Fig. 8; shows the structural formula of compound AMD 3530.
- Fig. 9; shows the structural formula of compound AMD 3517.
- Fig. 10; shows the structural formula of compound AMD 3544.
- Fig. 11; shows the structural formula of compound AMD 3543.
- 10 Fig. 12; shows the structural formula of compound AMD 3529.
- Fig. 13; shows the structural formula of compound AMD 7049.
- Fig. 14; shows the structural formula of compound AMD 7050.
- Fig. 15; shows the structural formula of compound AMD 7051.
- Fig. 16; shows the structural formula of compound AMD 7059.
- 15 Fig. 17; shows the structural formula of compound AMD 7063.
- Fig. 18; shows the structural formula of compound AMD 7058.
- Fig. 19; shows the structural formula of compound AMD 7032.
- Fig. 20; shows the structural formula of compound AMD 7048.
- Fig. 21; shows the structural formula of compound AMD 7060.
- 20 Fig. 22; shows the structural formula of compound AMD 7061.
- Fig. 23; shows the structural formula of compound AMD 3451.
- Fig. 24; shows the structural formula of compound AMD 3454.
- Fig. 25; shows the structural formula of compound AMD 3472.
- Fig. 26; shows the structural formula of compound AMD 3526.
- 25 Fig. 27; shows the structural formula of compound AMD 3100.
- Fig. 28; shows the structural formula of compound AMD 3484.
- Fig. 29, shows the cumulative incidence of collagen-induced arthritis in laboratory animals following treatment or immunization with compound AMD 3100.
- Fig. 30, shows the change in body weight of laboratory animals following treatment
- 30 with compound AMD 3100.
- Fig. 31, shows the structural formula of compound AMD 8630.

Fig. 32, shows the structural formula of compound AMD 7097.

Fig. 33, shows the structural formula of compound AMD 8631.

Fig. 34, shows the structural formula of compound AMD-Exp 1.

Fig. 35, shows the structural formula of compound AMD 7450.

5 Fig. 36, shows the structural formula of compound AMD-Exp 2.

Fig. 37, shows the structural formula of compound AMD 7463.

Fig. 38, shows the structural formula of compound AMD-Exp 3.

### DETAILED DESCRIPTION OF THE INVENTION

10 Terms as used herein are based upon their art recognized meaning unless otherwise indicated and should be clearly understood by the ordinary skilled artisan. The present invention will now be illustrated by the following preparative Examples.

#### General Procedure A

15 1-[1-Methylene-4-(bromomethylene)phenylene]-4,8,11-tris(diethoxyphosphoryl)-1,4,8,11-tetraazacyclotetradecane

To a stirred solution of 4,8,11-Tris(diethoxyphosphoryl)-1,4,8,11-tetraazacyclotetradecane (See Bridger *et al. J Med. Chem.* 1995, 38, 366-378) (6.1g, 0.01mol) and  $K_2CO_3$  (1.89g, 0.013mol) in  $CH_3CN$  (150ml) was added  $\alpha,\alpha'$ -dibromo-p-xylene (13.2g, 0.05mol) and the reaction mixture stirred at 70°C for 1 hour. The solution was cooled to room temperature and the solvent removed under reduced pressure. The residue was partitioned between brine (50ml) and  $CH_2Cl_2$  (100ml). The organic phase was separated, dried ( $Na_2SO_4$ ) and concentrated to a minimum volume. The solid was filtered off and the solvent evaporated under reduced pressure to give the crude product as a pale yellow oil. Purification by column chromatography in silica gel ( $CH_2Cl_2/CH_3OH$ , 25:1) gave 1-[1-methylene-4-(bromomethylene)phenylene]-4,8,11-tris(diethoxyphosphoryl)-1,4,8,11-tetraazacyclotetradecane (4.7g, 59%) as a pale yellow oil.  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.21-1.37 (m, 18H), 1.66-1.74 (m, 2H), 1.82-1.91 (m, 2H), 2.30-2.35 (m, 2H), 2.58-2.63 (m, 2H), 2.99-3.16 (m, 12H), 3.48 (s, 2H), 3.95-4.07 (m, 12H), 4.48 (s, 2H), 7.21-7.35 (4H).

### General Procedure B

Second alkylation of the bromobenzyl cyclam intermediate with an amine (see for example:

Bridger *et al. J. Med. Chem.* 1995, 38, 366-378)

To a solution of the appropriate amine (5.0 equiv.) in dry CH<sub>3</sub>CN (5mL) containing a  
5 suspension of K<sub>2</sub>CO<sub>3</sub> (1.5 equiv.) at 80°C was added dropwise with stirring a solution of 1-[1-  
methylene-4-(bromomethylene)phenylene]-4,8,11-tris(diethoxyphosphoryl)-1,4,8,11-  
tetraazacyclotetradecane (0.6mmol) in CH<sub>3</sub>CN (10ml) over 15-20 min. After stirring for a  
further 1 hour at 80°C the solution was concentrated to dryness and the residue was partitioned  
between CH<sub>2</sub>Cl<sub>2</sub> and water. The organic layer was separated and washed with water (3x) then  
10 dried (MgSO<sub>4</sub>) and evaporated. The crude residue was purified by column chromatography on  
silica gel eluting with 5-15% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to afford a viscous oil.

### General Procedure C

De-protection of the diethoxyphosphoramidate groups using HBr/HOAc at room temperature

15 (see for example: Bridger *et al. J. Med. Chem.* 1995, 38, 366-378)

To a stirred solution of the protected cyclam derivative from procedure B (0.1-0.5mmol)  
in acetic acid (3mL) was added 30% HBr in acetic acid (Aldrich, 5mL) and the solution was  
stirred at room temperature for 14 hours. The resulting precipitate was collected by filtration  
and washed with acetic acid then Et<sub>2</sub>O. The solid was then dissolved in H<sub>2</sub>O (3mL) and treated  
20 with charcoal (100mg) and the mixture was heated to 80°C for 30min. The hot solution was  
filtered through celite and the filtrate was concentrated to approximately 1mL after which acetic  
acid was added resulting in the immediate formation of a white precipitate. The white solid was  
collected by filtration and dried *in vacuo*.

The following compounds were prepared by these methods:

25

### Example 1

N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-methyl)pyridine  
hexahydrobromide (AMD 3465)

White solid: Mp 200-205°C (dec); <sup>1</sup>H NMR (D<sub>2</sub>O) δ 2.04 (m, 4H), 3.20-3.40 (m, 8H),  
30 3.40-3.60 (m, 8H), 4.34 (s, 2H), 4.38 (s, 2H), 4.51 (s, 2H), 7.50 (m, 4H), 7.75 (t, 1H, *J*=6.6 Hz),  
7.82 (d, 1H, *J*=7.9 Hz), 8.26 (t, 1H, *J*=7.9 Hz), 8.63 (d, 1H, *J*=5.3 Hz); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 18.30,

18.96, 37.04, 37.28, 37.40, 40.92, 41.13, 41.49, 44.26, 47.61, 48.01, 51.29, 58.88, 127.46, 127.75, 130.40, 131.05, 131.23, 131.47, 132.10, 132.44, 144.95, 145.81, 146.01; FAB MS  $m/z$  493 ( $M+H^{81}Br$ , 7), 491 ( $M+H^{79}Br$ , 7), 411 ( $M+H$ , 100).

Anal. ( $C_{24}H_{38}N_6 \cdot 6HBr$ ); Calc. C, 32.36; H, 4.98; N, 9.44; Br, 53.21. Found C, 32.20; H, 5.00; N, 9.30; Br, 53.10.

### Example 2

*N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-*N*-methyl-2-(aminomethyl)pyridine hexahydrobromide hydrate (AMD 3538)

10 White solid: Mp 220-225°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  2.06 (m, 4H), 2.76 (s, 3H), 3.20-3.65 (m, 16H), 4.47 (bs, 4H), 4.65 (s, 2H), 7.54 (bs, 4H), 7.80 (t, 1H), 7.87 (d, 1H), 8.28 (t, 1H), 8.68 (d, 1H);  $^{13}C$  NMR ( $D_2O$ )  $\delta$  18.14, 18.75, 18.89, 36.74, 37.04, 37.15, 37.62, 40.38, 40.72, 40.91, 41.28, 44.05, 47.50, 56.98, 58.88, 60.28, 127.60, 128.86, 130.78, 130.96, 132.16, 132.64, 144.91, 145.04, 146.12; FAB MS  $m/z$  507 ( $M+H^{81}Br$ , 27), 507 ( $M+H^{79}Br$ , 22), 425 ( $M+H$ , 100).

15 Anal. ( $C_{25}H_{40}N_6 \cdot 6HBr \cdot 1.5H_2O$ ); Calc. C, 32.04; H, 5.27; N, 8.97; Br, 51.16. Found C, 31.88; H, 5.30; N, 8.93; Br, 51.00.

### Example 3

*N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-amino-methylpyridine hexahydrobromide (AMD 3500)

20 White solid: mp 201-204°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  1.91-2.12 (m, 4H), 3.00-3.49 (m, 16H), 4.13 (s, 2H), 4.34 (s, 2H), 4.53 (s, 2H), 7.39-7.57 (m, 4H), 8.02 (d, 2H,  $J=6.3$  Hz), 8.74 (d, 2H,  $J=6.3$  Hz);  $^{13}C$  NMR ( $D_2O$ )  $\delta$  18.26, 18.88, 36.94, 37.29, 37.36, 40.89, 41.06, 41.44, 44.21, 47.61, 49.17, 51.43, 59.02, 127.84, 130.21, 131.64, 132.15, 132.45, 142.19, 151.67; FAB MS  $m/z$  493 ( $M+H^{81}Br$ , 8), 491 ( $M+H^{79}Br$ , 10), 411 ( $M+H$ , 83), 320 (37), 247 (58), 201 (100).

25 Anal. ( $C_{24}H_{38}N_6 \cdot 6HBr$ ); Calc. C, 32.17; H, 4.95; N, 9.34; Br, 53.50. Found C, 32.16; H, 5.03; N, 9.41; Br, 53.28.

Example 4*N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-3-(amino-methyl)pyridine  
hexahydrobromide (AMD 3499)

White solid: mp 198-202°C (dec); <sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.83-2.07 (m, 4H), 2.96-3.47 (m, 5 16H), 4.11 (s, 2H), 4.32 (s, 2H), 4.49 (s, 2H), 7.38-7.56 (m, 4H), 8.04 (t, 1H, *J*=6.4 Hz), 8.63 (d, 1H, *J*=8.3 Hz), 8.76 (d, 1H, *J*=5.6 Hz), 8.86 (s, 1H); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 18.23, 18.87, 36.92, 37.29 (2C), 40.88, 41.05, 41.43, 44.17, 47.22, 47.60, 51.18, 59.04, 128.29, 130.01, 131.49, 132.14, 132.66 (2C), 142.55, 142.76, 148.98; FAB MS *m/z* 493 (M+H<sup>81</sup>Br, 7), 491 (M+H<sup>79</sup>Br, 6), 411 (M+H, 100), 320 (33), 247 (24).

10        Anal. (C<sub>24</sub>H<sub>38</sub>N<sub>6</sub>·6HBr); Calc. C, 32.17; H, 4.95; N, 9.34; Br, 53.50. Found C, 32.08; H, 5.02; N, 9.25; Br, 53.28.

Example 5

15        *N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-(2-amino-methyl-5-methyl)pyrazine pentahydrobromide (AMD 3498)

White solid: mp 194-197°C (dec); <sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.93-2.12 (m, 4H), 2.42 (s, 3H), 3.25 (s, 8H), 3.48 (s, 8H), 4.28 (s, 2H), 4.30 (s, 2H), 4.33 (s, 2H), 7.44 (s, 4H), 8.33 (s, 1H), 8.46 (s, 1H); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 18.01, 18.72, 19.80, 36.66, 37.05, 37.13, 40.70, 40.89, 41.27, 43.99, 47.47, 48.14, 50.61, 59.06, 129.97, 131.43, 132.04, 132.99, 140.93, 144.98, 146.49, 20 153.51; FAB MS *m/z* 509 (M+H<sup>81</sup>Br, 17), 507 (M+H<sup>79</sup>Br, 15), 426 (M+H, 100), 320 (21), 247 (20).

      Anal. (C<sub>24</sub>H<sub>39</sub>N<sub>7</sub>·5.5HBr); Calc. C, 33.10; H, 5.15; N, 11.26; Br, 50.47. Found C, 32.80; H, 5.41; N, 11.00; Br, 50.58.

25

Example 6

*N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-ethyl)pyridine  
hexahydrobromide (AMD 3497)

White solid: mp 195-198°C (dec); <sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.98-2.17 (m, 4H), 3.20-3.38 (m, 8H), 3.38-3.63 (m, 12H), 4.27 (s, 2H), 4.39 (s, 2H), 7.50 (s, 4H), 7.80-7.89 (m, 2H), 8.42 (m, 30 1H), 8.58 (d, 1H, *J*=5.8 Hz); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 18.51, 19.14, 29.85, 37.56 (3C), 41.21, 41.41,



41.82, 44.57, 45.27, 47.83, 51.10, 58.74, 126.35, 127.93, 130.66, 131.27, 131.99, 132.69, 141.89, 147.79, 150.91; FAB MS  $m/z$  507 ( $M+H^{81}Br$ , 40), 505 ( $M+H^{79}Br$ , 34), 425 ( $M+H$ , 100).

Anal. ( $C_{25}H_{40}N_6 \cdot 6HBr$ ); Calc. C, 32.99; H, 5.09; N, 9.23; Br, 52.67. Found C, 32.79; H, 5.34; N, 9.11; Br, 52.45.

5

#### Example 7

##### *N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-methyl)thiophene pentahydrobromide (AMD 3516)

White solid: mp 245-248°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  1.87-2.12 (m, 4H), 3.02-3.51 (m, 16H), 4.17 (s, 4H), 4.38 (s, 2H), 6.97 (t, 1H,  $J=3.9$  Hz), 7.13 (d, 1H,  $J=3.1$  Hz), 7.41 (s, 5H);  $^{13}C$  NMR ( $D_2O$ )  $\delta$  18.80, 19.52, 38.03, (3C), 41.59 (2C), 42.21, 44.89 (2C), 48.15, 49.83, 58.52, 128.13, 129.12, 131.15, 131.47, 131.50, 131.90, 132.42, 132.87; FAB MS  $m/z$  498 ( $M+H^{81}Br$ , 11), 496 ( $M+H^{79}Br$ , 9), 416 ( $M+H$ , 53), 218 (100), 201 (64).

Anal. ( $C_{23}H_{37}N_5S \cdot 5HBr$ ); Calc. C, 33.68; H, 5.16; N, 8.54; Br, 48.71. Found C, 33.85; H, 5.22; N, 8.50; Br, 48.52.

15

#### Example 8

##### *N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-ethyl)mercaptan pentahydrobromide dihydrate (AMD 3530)

White solid: mp 234-236°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  1.75-2.05 (m, 4H), 2.75-3.45 (m, 20H), 4.05 (s, 2H), 4.15 (s, 2H), 7.35 (s, 4H); FAB MS  $m/z$  462 ( $MH+H^{81}Br$ , 15), 460 ( $MH+H^{79}Br$ , 15), 380 ( $M+H$ , 100), 300 (64), 279 (47), 239 (49).

Anal. ( $C_{20}H_{37}N_5S \cdot 5HBr \cdot 2H_2O \cdot 0.5HOAc$ ) requires C, 29.67; H, 5.69; N, 8.24; Br, 46.99. Found C, 29.31; H, 5.72; N, 8.25; Br, 46.64.

25

#### Example 9

##### *N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-amino-benzylamine pentahydrobromide (AMD 3517)

White solid: mp 203-206°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  1.85-2.13 (m, 4H), 3.02-3.58 (m, 16H), 4.23 (s, 2H), 4.31 (s, 4H), 7.23-7.54 (m, 8H);  $^{13}C$  NMR ( $D_2O$ )  $\delta$  18.03, 19.29, 37.78 (3C), 41.37 (2C), 42.00, 44.82, 46.25, 47.96, 51.16, 58.68, 124.04, 124.40, 129.40, 130.75, 131.21

30

(2C), 131.88, 131.96, 132.46, 132.83; FAB MS  $m/z$  507 ( $M+H^{81}Br$ , 15), 505 ( $M+H^{79}Br$ , 18), 425 ( $M+H$ , 100), 320 (30), 201 (51).

Anal. ( $C_{25}H_{40}N_6 \cdot 5.75HBr \cdot 0.5H_2O$ ). Calc. C, 33.42; H, 5.19; N, 9.35; Br, 51.14. Found C, 33.69; H, 5.35; N, 9.00; Br, 51.13.

5

#### Example 10

*N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-amino-benzylamine  
hexahydrobromide (AMD 3544)

Yellow solid: mp 120-125°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  1.8-2.0 (m, 4H), 2.9-3.4 (m, 10 16H), 4.1 (s, 2H), 4.18 (s, 4H), 7.2-7.5 (m, 8H);  $^{13}C$  NMR ( $D_2O$ )  $\delta$  18.86, 19.57, 38.14, 41.76, 43.74, 45.14, 48.24, 50.14, 50.42, 51.49, 58.38, 124.13, 131.13, 131.30, 131.83, 131.92, 131.96, 132.67; FAB MS  $m/z$  507 ( $M+H^{81}Br$ , 5), 505 ( $M+H^{79}Br$ , 5), 425 ( $M+H$ , 45), 201 (47), 155 (75), 106 (100).

Anal. ( $C_{25}H_{40}N_6 \cdot 6HBr \cdot HOAc$ ) requires C, 33.43; H, 5.19; N, 8.66; Br, 49.42; O, 3.30.  
15 Found C, 33.42; H, 5.49; N, 8.62; Br, 49.23.

#### Example 11

*N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-(amino-ethyl)imidazole  
hexahydrobromide (AMD 3543)

20 Off white solid: mp 135-140°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  1.75 (m, 2H), 1.90 (m, 2H), 2.70-3.27 (m, 20H), 3.77 (s, 2H), 4.14 (s, 2H), 7.18 (s, 1H), 7.25 (d, 2H,  $J=7.97$  Hz), 7.37 (d, 2H,  $J=7.97$  Hz), 8.48 (s, 1H); FAB MS  $m/z$  496 ( $M+H^{81}Br$ , 5), 494 ( $M+H^{79}Br$ , 5), 414 ( $M+H$ , 17), 201 (15).

Anal. ( $C_{23}H_{39}N_7 \cdot 6HBr$ ) requires C, 30.73; H, 5.04; N, 10.91; Br, 53.32. Found C, 30.39;  
25 H, 5.41; N, 10.41; Br, 53.66.

#### Example 12

*N*-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-benzylamine  
pentahydrobromide (AMD 3529)

30 Off white solid: mp 245-250°C (dec);  $^1H$  NMR ( $D_2O$ )  $\delta$  1.9-2.1 (m, 4H), 3.2-3.6 (m, 16H), 4.12 (s, 2H), 4.15 (s, 2H), 4.36 (s, 2H), 7.30 (s, 5H), 7.41 (d, 2H,  $J=8.3$  Hz) 7.46 (d, 2H,

$J=8.3$  Hz);  $^{13}\text{C}$  NMR ( $\text{D}_2\text{O}$ )  $\delta$  18.43, 19.06, 37.29, 37.46, 37.63, 41.09, 41.32, 41.68, 44.46, 47.74, 50.18, 51.00, 58.79, 129.53, 129.97, 130.18, 130.35, 130.68, 131.18, 131.92, 133.14; FAB MS  $m/z$  492 ( $\text{M}+\text{H}^{81}\text{Br}$ , 13), 490 ( $\text{M}+\text{H}^{79}\text{Br}$ , 13), 410 ( $\text{M}+\text{H}$ , 100), 201 (36).

Anal. ( $\text{C}_{25}\text{H}_{39}\text{N}_5 \cdot 5\text{HBr}$ ); requires C, 36.88; H, 5.45; N, 8.60; Br, 49.07. Found C, 36.79;  
5 H, 5.56; N, 8.48; Br, 48.79.

The compounds of the invention were tested in a screen by the MTT method (*J. Virol. Methods* 120:309-321 (1988)). MT-4 cells ( $2.5 \times 10^4/\text{well}$ ) were challenged with HIV-1 (HTLV-III<sub>B</sub>) or HIV-2 (LAV-2 ROD) at a concentration of 100 CCID<sub>50</sub> and incubated in the presence of various concentrations of the test compounds, which were added immediately after  
10 challenge with the virus. After 5 days culture at 37°C in a CO<sub>2</sub> incubator, the number of viable cells was assessed by the MTT (tetrazolium) method. Antiviral activity and cytotoxicity of the compounds are expressed in Table 1 below as EC<sub>50</sub> ( $\mu\text{g}/\text{ml}$ ) and CC<sub>50</sub> ( $\mu\text{g}/\text{ml}$ ), respectively. The potential therapeutic usefulness was assessed by calculating a Selectivity Index (SI) corresponding to the ratio of CC<sub>50</sub> to EC<sub>50</sub>.

15

Table 1				
Anti-HIV activity data				
Compound	CC <sub>50</sub> ( $\mu\text{g}/\text{mL}$ )	EC <sub>50</sub> ( $\mu\text{g}/\text{mL}$ )		SI
		HIV-1 (IIIB)	HIV-2	HIV-1
1 AMD3465	>250	0.008	0.032	$3 \times 10^4$
2 AMD3538	209	0.1	6.7	$2.0 \times 10^3$
3 AMD3500	>250	0.6	10.3	417
4 AMD3499	>250	1.8	28.5	138
5 AMD3498	>250	0.2	7.1	$1.2 \times 10^3$
6 AMD3497	>250	1.8	3.8	138
7 AMD3516	158	0.7	9.8	225
8 AMD3530	175	0.5	2.0	350
9 AMD3517	153	0.8	10.6	191
10 AMD3544	222	0.7	3.7	317
11 AMD3543	239	0.2	1.0	$1 \times 10^3$
12 AMD3529	130	0.4	2.6	325

In this field of study, it is considered that any compound exhibiting a Selectivity Index of greater than 100 has the considerable potential for further study. HIV is one of the most challenging viruses to combat, and the results given above provide an indication of activity  
20 against other retroviruses and against other viruses in general.

Example 13

N-[4-(1,4,7-Triazacyclotetra-decanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine  
(AMD 7049)

5

N,N'-Bis(2-Nitrobenzenesulfonyl)-1,7-heptanediamine

To a stirred solution of 1,7-heptanediamine (5.01 g, 38.5 mmol) and Et<sub>3</sub>N (13.5 mL, 96.9 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (70 mL) was added a solution of 2-nitrobenzenesulfonyl chloride (18.80 g, 84.83 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL). The mixture was stirred at room temperature under  
10 nitrogen for 72 h and then concentrated *in vacuo*. The residue was stirred in diethyl ether (100 mL), and the precipitate was collected by filtration and washed with H<sub>2</sub>O (300 mL) followed by diethyl ether (300 mL) to give a gray solid (18.5 g, 96%): <sup>1</sup>H NMR (DMF-*d*<sub>7</sub>)  $\delta$  1.21 (m, 6H), 1.49 (m, 4H), 3.04 (m, 4H), 7.87 (m, 2H), 7.95 (m, 4H), 8.04 (m, 2H), 8.15 (m, 2H).

15

General Procedure D

4-Diethoxyphosphoryl-1,7-bis(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetradecane

To a stirred solution of N,N'-bis(2-nitrobenzenesulfonyl)-1,7-heptanediamine (9.00 g, 18.0 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (17.8 g, 54.6 mmol) in DMF (500 mL) under nitrogen maintained at 80 °C was added dropwise a solution of N-(diethoxyphosphoryl)-O,O'-bis(2-methylsulfonyl)di-ethanolamine (Bridger et al., J. Med. Chem. 1995, 38, 366-378) (7.95 g, 20.0  
20 mmol) in DMF (50 mL) over 8 h. Heating was continued for a further 17 h and the mixture was then allowed to cool and concentrated *in vacuo*. The residue was partitioned between CHCl<sub>3</sub> (140 mL) and H<sub>2</sub>O (80 mL) and the aqueous layer was separated and extracted with CHCl<sub>3</sub> (3 × 40 mL). The combined organic extracts were dried (MgSO<sub>4</sub>) and concentrated *in vacuo* and the  
25 residue was purified by column chromatography on silica gel (ethylacetate) to give the desired macrocycle as a yellow crystalline solid (2.85 g, contaminated with DMF).

To remove the unwanted DMF impurity, the residue was dissolved in EtOAc (75 mL), and the solution was washed sequentially with 5% NaHCO<sub>3</sub> (2 × 10 mL) and brine (5 × 10 mL), dried (MgSO<sub>4</sub>) and evaporated to give a yellow amorphous solid (2.52 g, 20%): <sup>1</sup>H NMR  
30 (CDCl<sub>3</sub>)  $\delta$  1.32 (t, 6H, *J* = 7.1 Hz), 1.51 (m, 6H), 1.61 (m, 4H), 3.33 (m, 12H), 4.03 (m, 4H), 7.61 (m, 2H), 7.71 (m, 4H), 8.03 (m, 2H).

### General Procedure E

#### Synthesis of 1,7-Bis(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetradecane

To a stirred suspension of the macrocycle from above (1.88 g, 2.66 mmol) in acetic acid (5 mL) was added a freshly prepared solution of saturated HBr(g) in acetic acid (20 mL) and the resulting homogeneous solution was stirred at room temperature for a further 22 h. Addition of diethyl ether (250 mL) to the reaction mixture gave a precipitate that was allowed to settle to the bottom of the flask and the supernatant solution was decanted. The precipitate was washed with ether by decantation (repeated 3×) and the residue was then partitioned between CH<sub>2</sub>Cl<sub>2</sub> (40 mL) and 1N aqueous NaOH (25 mL). The separated aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 × 20 mL) and the combined organic extracts were washed with brine (20 mL), then dried (MgSO<sub>4</sub>) and concentrated *in vacuo* to give a yellow amorphous solid (1.23 g, 81%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.46-1.67 (m, 10H), 2.90 (m, 4H), 3.34 (m, 8H), 7.61 (m, 2H), 7.70 (m, 4H), 7.97 (m, 2H).

#### 4-Bromomethylbenzyl alcohol

To a solution of methyl 4-bromomethylbenzoate (5.73 g, 25 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (150 mL) cooled to -78 °C with stirring under nitrogen was added dropwise a solution of DIBAL-H (82.5 mL, 1.0 M solution in THF). Stirring was continued for 1.5 h at -78 °C, and the reaction mixture was then allowed to warm to 0 °C and quenched with H<sub>2</sub>O. The organic layer was separated and the aqueous was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 100 mL). The combined organic extracts were dried (MgSO<sub>4</sub>) and evaporated to give the desired alcohol (5.0 g, 100%) as a white solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.84 (br, 1H), 4.49 (s, 2H), 4.67 (s, 2H), 7.33 (d, 2H, *J* = 8.2 Hz), 7.38 (d, 2H, *J* = 8.2 Hz).

#### N-(2-Nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

A solution of 2-nitrobenzenesulfonylchloride (16.62 g, 0.075 mol) in dry CH<sub>2</sub>Cl<sub>2</sub> (120 mL) was added dropwise via cannula to a stirred solution of 2-(aminomethyl)pyridine (5.41 g, 0.05 mol) and Et<sub>3</sub>N (13.9 mL, 0.10 mol) in dry CH<sub>2</sub>Cl<sub>2</sub> (150 mL) under nitrogen. The reaction mixture was stirred for three hours at room temperature, and then quenched with water (20 mL). The aqueous layer was separated and extracted with EtOAc (5 x 80 mL). The combined organic extracts were dried (MgSO<sub>4</sub>) and evaporated to small volume to give a white precipitate

which was collected by filtration and washed with cold  $\text{CH}_2\text{Cl}_2$  to give the desired product (11.37 g, 78%) as a white solid:  $^1\text{H}$  NMR (Acetone- $d_6$ )  $\delta$  4.46 (s, 2H), 7.19 (dd, 1H,  $J = 7.4, 4.5$  Hz), 7.25-7.35 (br s, 1H), 7.39 (d, 1H,  $J = 7.7$  Hz), 7.68 (ddd, 1H,  $J = 7.7, 7.5, 1.8$  Hz), 7.76-7.88 (m, 2H), 7.94 (dd, 1H,  $J = 7.7, 1.5$  Hz), 8.04 (dd, 1H,  $J = 7.5, 1.8$  Hz), 8.38 (d, 1H,  $J = 4.5$  Hz).

N-[1-Methylene-4-(hydroxymethylene)phenylene]-N-(2-Nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

A mixture of N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (5.87 g, 20 mmol), 4-bromomethylbenzyl alcohol (4.02 g, 20 mmol) and  $\text{K}_2\text{CO}_3$  (5.53 g, 40 mmol) in dry  $\text{CH}_3\text{CN}$  (150 mL) were heated at 60 °C for 4 h with stirring under nitrogen. The mixture was then allowed to cool to room temperature, the solvent evaporated and the residue was partitioned between water and  $\text{CH}_2\text{Cl}_2$ . The separated aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$ , and the combined organic extracts were dried ( $\text{MgSO}_4$ ) and evaporated. The residue was suspended in ethyl acetate/hexane (1:1) and collected by filtration to give the desired product (6.87 g, 83%) as a white solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.78 (t, 1H,  $J = 5.8$  Hz), 4.58 (s, 2H), 4.60 (s, 2H), 4.64 (d, 2H,  $J = 5.8$  Hz), 7.13-7.26 (m, 6H), 7.54-7.59 (m, 2H), 7.66-7.68 (m, 2H), 7.98 (d, 1H,  $J = 7.4$  Hz), 8.40 (d, 1H,  $J = 3.8$  Hz).

N-[1-Methylene-4-(chloromethylene)phenylene]-N-(2-Nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

To a stirred solution of the alcohol from above (1.91 g, 4.62 mmol) and  $\text{Et}_3\text{N}$  (2.0 mL, 14 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL) cooled in an ice bath under nitrogen, was added methanesulfonyl chloride (0.73 mL, 9.4 mmol) and the reaction mixture was then heated to reflux for a further 6 h. The solution was diluted with  $\text{CH}_2\text{Cl}_2$  (60 mL) and washed with 10% aqueous HCl (2  $\times$  20 mL), 5% aqueous  $\text{NaHCO}_3$  (20 mL), and  $\text{H}_2\text{O}$  (25 mL) then dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo* to give an orange oil (1.95 g, 98%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  4.52 (s, 2H), 4.60 (s, 4H), 7.12-7.26 (m, 6H), 7.55 (m, 2H), 7.67 (d, 2H,  $J = 4.0$  Hz), 7.94 (d, 1H,  $J = 8.0$  Hz), 8.41 (d, 1H,  $J = 4.8$  Hz). This was used without further purification.

General Procedure F

N-[4-[1,7-Bis(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetra-decanyl]-1,4-

phenylenebis(methylene)]-N-(2-Nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

A mixture of 1,7-bis(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetradecane (1.1 g, 1.9 mmol), the chloride from above (0.98 g, 2.3 mmol) and  $K_2CO_3$  (0.85 g, 6.2 mmol) were heated to reflux in  $CH_3CN$  (30 mL) under nitrogen for 62 h. The solvent was evaporated in vacuo and the residue was partitioned between  $CH_2Cl_2$  (100 mL) and brine (70 mL). The aqueous phase was separated and extracted with  $CH_2Cl_2$  (40 mL) and the combined organic extracts were dried ( $MgSO_4$ ) and concentrated in vacuo. The residue was purified by column chromatography on silica gel (3% MeOH/ $CH_2Cl_2$ ) and the evaporated fractions containing the desired product were subjected to a second column purification on silica gel (ethyl acetate) to give a pale yellow amorphous solid (940 mg, 49%):  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.44 (br s, 6H), 1.60 (br s, 4H), 2.75 (m, 4H), 3.23-3.33 (m, 8H), 3.59 (s, 2H), 4.58 (s, 2H), 4.59 (s, 2H), 7.08-7.20 (m, 6H), 7.55-7.70 (m, 10H), 7.82 (dd, 2H,  $J = 7.6, 1.6$  Hz), 7.99 (d, 1H,  $J = 7.8$  Hz), 8.40 (d, 1H,  $J = 4.7$  Hz).

N-[4-(1,4,7-Triazacyclotetra-decanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine Pentahydrobromide Dihydrate

The intermediate from above (870 mg, 0.90 mmol),  $K_2CO_3$  (1.15 g, 8.32 mmol), and thiophenol (0.33 mL, 3.2 mmol) were stirred in DMF (12 mL) for 7.5 h at room temperature. The mixture was concentrated *in vacuo* and the residue was partitioned between  $CH_2Cl_2$  (30 mL) and  $H_2O$  (15 mL). The organic phase was separated, washed with 5%  $NaHCO_3$  (10 mL) then  $H_2O$  (10 mL) then dried ( $MgSO_4$ ) and concentrated *in vacuo*. The yellow residue was purified by column chromatography on basic alumina ( $CH_2Cl_2$ , 1% MeOH/ $CH_2Cl_2$ , and 10% MeOH/ $CH_2Cl_2$ ) to give the free base as a yellow oil (134 mg, 36%):  $^1H$  NMR ( $CDCl_3$ )  $\delta$  1.48 (br s, 6H), 1.60 (br s, 4H), 2.61 (m, 12H), 3.56 (s, 2H), 3.83 (s, 2H), 3.92 (s, 2H), 7.16 (m, 1H), 7.24 (m, 2H), 7.32 (m, 3H), 7.79 (m, 1H), 8.56 (d, 1H,  $J = 4.7$  Hz).

The free base (134 mg, 0.33 mmol) was dissolved in EtOH (4 mL) and a freshly prepared solution of saturated HBr(g) in EtOH (9 mL) was added, giving a white precipitate. The mixture was stirred for 5 min and diethyl ether (15 mL) was added. The precipitate was allowed to settle to the bottom of the flask and the supernatant solution was decanted. The solid was then dissolved in MeOH (5 mL) and re-precipitated with a large volume of ether, washed with ether by decantation (15x) and finally, the last traces of ether were removed by evaporation at reduced pressure (room temperature). Drying the solid *in vacuo* at 40 °C for 16 h, gave the

desired product as a white solid (178 mg, 63%):  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  1.44 (br s, 6H), 1.75 (br s, 4H), 3.04 (br s, 8H), 3.37 (m, 4H), 4.06 (br s, 2H), 4.31 (s, 2H), 4.38 (s, 2H), 7.52-7.68 (m, 6H), 8.01 (m, 1H), 8.70 (d, 1H,  $J = 5.0$  Hz); FAB-MS  $m/z$  492 (MH +  $\text{H}^{81}\text{Br}$ ), 490 (MH +  $\text{H}^{79}\text{Br}$ ), 410 (M + H). Anal. Calcd for  $\text{C}_{25}\text{H}_{39}\text{N}_5 \cdot 5\text{HBr} \cdot 0.1\text{Et}_2\text{O} \cdot 2.3\text{H}_2\text{O}$ : C, 35.35; H, 5.79; N, 8.11; Br, 46.29. Found: C, 35.55; H, 5.70; N, 8.18; Br, 46.17.

#### Example 14

##### N-[7-(4,7,10,17-Tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7050)

10

2,6-Bis(2-aminoethyl)pyridine was prepared as described in Bridger et al. U.S. Pat. No. 5,698,546, which is hereby incorporated in its entirety by reference herein.

##### 2,6-Bis[N-(2-nitrobenzenesulfonyl)-2-aminoethyl]pyridine

To a stirred solution of 2,6-Bis(2-aminoethyl)pyridine (2.7 g, 16 mmol) and  $\text{Et}_3\text{N}$  (5.7 mL, 41 mmol) in  $\text{CH}_2\text{Cl}_2$  (35 mL) was added 2-nitrobenzenesulfonyl chloride (8.01 g, 36.1 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL) and the mixture was stirred at room temperature under nitrogen for 42 h. The mixture was washed with brine (25 mL) and the organic phase was dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo*. The brown residue was purified by column chromatography on silica gel (50% then 60% THF/hexane) to give a pale yellow solid (5.2 g, 59%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  3.01 (m, 4H), 3.52 (m, 4H), 6.38 (m, 2H), 6.94 (d, 2H,  $J = 7.7$ , Hz), 7.47 (t, 1H,  $J = 7.7$  Hz), 7.72 (m, 4H), 7.82 (m, 2H), 8.13 (m, 2H).

20

##### 7-Diethoxyphosphoryl-4,10-Bis(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene

Using General Procedure D: Reaction of 2,6-bis[N-(2-nitrobenzenesulfonyl)-2-aminoethyl]pyridine (5.2 g, 9.7 mmol) and N-(diethoxyphosphoryl)-O,O'-bis(2-methylsulfonyl)di-ethanolamine (4.25 g, 10.7 mmol) followed by silica gel column purification (60% then 90% THF/hexane) of the reaction products gave the title compound as a yellow amorphous solid (1.48 g, 21%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.23 (t, 6H,  $J = 7.1$  Hz), 2.60 (m, 4H), 2.98-3.08 (m, 8H), 3.84-3.94 (m, 8H), 7.11 (d, 2H,  $J = 7.6$  Hz), 7.56-7.74 (m, 7H), 8.07 (m, 2H).

30

##### 4,10-Bis(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-



1(17),13,15-triene

Using General Procedure E: Reaction of 7-diethoxyphosphoryl-4,10-bis(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene (1.04 g, 1.4 mmol) gave the title compound as a yellow amorphous solid (744 mg, 88%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.81 (m, 4H), 3.08 (m, 4H), 3.33 (m, 4H), 3.88 (m, 4H), 7.07 (d, 2H, J = 7.7 Hz), 7.54-7.71 (m, 7H), 8.02 (m, 2H).

N-[7-[4,10-Bis(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

Using General Procedure F: Reaction of 4,10-bis(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene (740 mg, 1.2 mmol) and N-[1-methylene-4-(chloromethylene)phenylene]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (610 mg, 1.4 mmol) followed by silica gel column purification (50% then 80% THF/hexane) of the reaction products gave the title compound as a yellow amorphous solid (648 mg, 54%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 2.26 (m, 4H), 3.03 (m, 8H), 3.37 (s, 2H), 3.94 (m, 4H), 4.56 (s, 2H), 4.57 (s, 2H), 6.95-7.17 (m, 8H), 7.52-7.72 (m, 11H), 7.85 (m, 2H), 7.98 (d, 1H, J = 7.7 Hz), 8.39 (d, 1H, J = 4.8 Hz).

General Procedure G

N-[7-(4,7,10,17-Tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine Hexahydrobromide Trihydrate

To a solution of N-[7-[4,10-bis(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (640 mg, 0.64 mmol) in DMF (9 mL) containing K<sub>2</sub>CO<sub>3</sub> (806 mg, 5.83 mmol) was added thiophenol (0.24 mL, 2.3 mmol) and the mixture was stirred at room temperature for 2 h. The mixture was concentrated in vacuo and the residue partitioned between ethyl acetate (30 mL) and water (10 mL). The organic phase was separated and extracted with 5% NaHCO<sub>3</sub> (3 x 5 mL) then brine (5 mL). The combined aqueous phases were extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 10 mL). The combined organic extracts were dried (MgSO<sub>4</sub>) and evaporated and the residue was purified by column chromatography on alumina (CH<sub>2</sub>Cl<sub>2</sub> followed by 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give the free base of the title compound as a

yellow oil (83 mg, 29%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.57 (m, 8H), 3.01 (s, 8H), 3.36 (s, 2H), 3.78 (s, 2H), 3.92 (s, 2H), 6.64 (d, 2H,  $J = 8.0$  Hz), 7.07 (m, 4H), 7.18 (m, 1H), 7.33 (d, 1H,  $J = 7.7$  Hz), 7.67 (m, 2H), 8.58 (d, 1H,  $J = 4.8$  Hz).

The free base (74 mg, 0.17 mmol) was dissolved in MeOH (3 mL) and a freshly prepared solution of saturated HBr(g) in MeOH (7 mL) was added giving a white precipitate. The mixture was stirred for 5 min and diethyl ether was added (10 mL), the solid was allowed to settle to the bottom of the flask and the supernatant solution decanted. The solid was washed by decantation with MeOH (5 x 5 mL) then ether (10 x 5 mL) and the last traces of ether were removed by evaporation *in vacuo* followed by drying *in vacuo* at 40 °C for 17.5 h to give the title compound as a white solid (153 mg, 93%):  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  2.81 (br s, 4H), 3.28 (m, 8H), 3.61 (br s, 4H), 3.85 (s, 2H), 4.27 (s, 2H), 4.36 (s, 2H), 7.29 (d, 2H,  $J = 7.7$  Hz), 7.36 (d, 2H,  $J = 7.7$  Hz), 7.53 (m, 3H), 7.63 (d, 1H,  $J = 7.7$  Hz), 7.80 (t, 1H,  $J = 7.7$  Hz), 7.99 (m, 1H), 8.69 (d, 1H,  $J = 5.3$  Hz); FAB-MS  $m/z$  527 ( $\text{MH} + \text{H}^{81}\text{Br}$ ), 525 ( $\text{MH} + \text{H}^{79}\text{Br}$ ), 445 ( $\text{M} + \text{H}$ ). Anal. Calcd for  $\text{C}_{27}\text{H}_{36}\text{N}_6 \cdot 6\text{HBr} \cdot 3\text{H}_2\text{O}$ : C, 32.95; H, 4.92; N, 8.54; Br, 48.72. Found: C, 32.75; H, 4.89; N, 8.39; Br, 48.61.

#### Example 15

N-[7-(4,7,10-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD7051)

#### 1,3-Phenylenebis(ethylene)diamine

To a solution of 1,3-phenylenediacetonitrile (9.37 g, 60 mmol) in  $\text{CH}_3\text{OH}$  (saturated with  $\text{NH}_3$ , 150 mL) was added Raney-Ni (ca. 20 g, previously washed with  $\text{CH}_3\text{OH}$  several times) and the mixture was hydrogenated at 45 psi on a Parr apparatus for 48 h. The reaction mixture was filtered through celite and the filtrate evaporated to give the crude product (9.45 g, 96%) as a light green oil:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  0.80-1.50 (br s, 4H), 2.70-2.76 (m, 4H), 2.94-2.99 (m, 4H), 7.01-7.07 (m, 3H), 7.18-7.26 (m, 1H). This was used in the next step without further purification.

#### N,N'-Bis(2-Nitrobenzenesulfonyl)-1,3-Phenylenebis(ethylene)diamine

A solution of 2-nitrobenzenesulfonylchloride (19.94 g, 0.090 mol) in dry  $\text{CH}_2\text{Cl}_2$  (70 mL) was added dropwise via cannula to a stirred solution of 1,3-phenylenebis(ethylene)diamine

(4.92 g, 0.030 mol) and Et<sub>3</sub>N (16.7 mL, 0.12 mol) in dry CH<sub>2</sub>Cl<sub>2</sub> (80 mL) under nitrogen. The reaction mixture was stirred overnight at room temperature, and then quenched with water (20 mL). The precipitate was collected by filtration and washed with H<sub>2</sub>O, CH<sub>3</sub>OH, and Et<sub>2</sub>O to give the desired product (9.22 g, 58%) as a white solid: <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.66 (t, 4H, J = 7.7 Hz), 3.08-3.18 (br s, 4H), 6.94 (d, 2H, J = 6.4 Hz), 6.98 (s, 1H), 7.12 (dd, 1H, J = 6.4, 6.4 Hz), 7.78-7.84 (br m, 4H), 7.90-7.64 (br m, 4H), 8.16 (br s, 2H).

7-Diethoxyphosphoryl-4,10-bis(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-triene

10 Using General Procedure D: Reaction of N,N'-bis(2-nitrobenzenesulfonyl)-1,3-phenylenebis(ethylene)diamine (8.74 g, 16.4 mmol) with N-(diethoxyphosphoryl)-O,O'-bis(2-methylsulfonyl)di-ethanolamine (6.50 g, 16.4 mmol) followed by silica gel column purification of the reaction products (1:15:35 CH<sub>3</sub>OH-Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound (4.03 g, 33%) as a yellow foam: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.21 (t, 6H, J = 6.4 Hz), 2.39-2.46 (br m, 4H), 2.83-2.97 (br m, 8H), 3.68-3.72 (m, 4H), 3.80-3.92 (m, 4H), 7.16 (d, 2H, J = 6.5 Hz), 7.18 (s, 1H), 7.24 (dd, 1H, J = 6.5, 6.5 Hz), 7.60-7.74 (m, 6H), 8.04-8.08 (m, 2H).

4,10-Bis(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-triene

20 Using General Procedure E: Reaction of 7-diethoxyphosphoryl-4,10-bis(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-triene (1.27 g, 1.72 mmol) followed by silica gel column purification of the reaction products (1:15:25 CH<sub>3</sub>OH-EtOAc-CH<sub>2</sub>Cl<sub>2</sub> then 20% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound (574 mg, 57%) as a light yellow foam: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.42-1.50 (br, 1H), 2.01 (t, 4H, J = 5.4 Hz), 2.90-3.10 (br m, 4H), 3.08 (t, 4H, J = 5.4 Hz), 3.56-3.60 (br m, 4H), 7.16 (d, 2H, J = 6.8 Hz), 7.31 (dd, 1H, J = 6.8, 6.8 Hz), 7.36 (s, 1H), 7.61-7.63 (m, 2H), 7.70-7.73 (m, 4H), 8.01-8.04 (m, 2H).

N-[7-[4,10-Bis(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-Nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

30 Using General Procedure F: Reaction of 4,10-bis(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-triene (420 mg, 0.7 mmol) with N-[1-methylene-4-(chloromethylene)phenylene]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (302 mg,

0.7 mmol) followed by silica gel column purification of the reaction products (1:3 Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound (491 mg, 70%) as a pale yellow solid: <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.97-2.02 (br m, 4H), 2.73-2.78 (br m, 4H), 2.90-2.94 (br m, 4H), 3.32 (s, 2H), 3.64-3.67 (br m, 4H), 4.55 (s, 2H), 4.58 (s, 2H), 6.93 (d, 2H, *J* = 8.0 Hz), 7.04 (d, 2H, *J* = 8.0 Hz), 7.09-7.16 (br m, 4H), 7.23 (s, 1H), 7.29 (dd, 1H, *J* = 7.9, 7.9 Hz), 7.51-7.72 (m, 10H), 7.80-7.83 (m, 2H), 7.98 (d, 1H, *J* = 7.8 Hz), 8.39 (m, 1H).

N-[7-(4,7,10-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine Pentahydrobromide Dihydrate

Using General Procedure G: Reaction of N-[7-[4,10-bis(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (380 mg, 0.38 mmol) followed by basic alumina column purification of the reaction products (1:20 CH<sub>3</sub>OH-CH<sub>2</sub>Cl<sub>2</sub>) gave the free base of the title compound.

Conversion of the free base to the hydrobromide salt using a saturated solution of HBr(g) in CH<sub>3</sub>OH followed by drying *in vacuo* overnight, gave the title compound (110 mg, 34% overall) as a white solid: <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 2.80-2.88 (br s, 4H), 3.02-3.06 (br s, 4H), 3.10-3.16 (br s, 4H), 3.38-3.44 (br s, 4H), 3.80-3.86 (br s, 2H), 4.25-4.30 (br s, 2H), 4.33-4.37 (br s, 2H), 7.27-7.32 (br m, 4H), 7.42-7.63 (br m, 6H), 7.96 (dd, 1H, *J* = 7.7, 7.7 Hz), 8.10-8.30 (br s, 3H), 8.69 (d, 1H, *J* = 4.9 Hz), 9.45-9.62 (br s, 2H); FAB-MS *m/z* 526 (MH + H<sup>81</sup>Br), 524 (MH + H<sup>79</sup>Br), 444 (M + H, 100); Anal. Calcd for C<sub>28</sub>H<sub>42</sub>N<sub>5</sub>Br<sub>5</sub>•2H<sub>2</sub>O: C, 38.03; H, 5.24; N, 7.92; Br, 45.18. Found: C, 38.37; H, 5.28; N, 7.76; Br, 45.36.

Example 16

N-[1-(1,4,7-Triazacyclotetra-decanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7059)

25

General Procedure H

4-Diethoxyphosphoryl-7-(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetradecane

To a stirred solution of 4-diethoxyphosphoryl-1,7-bis(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetradecane (1.32 g, 1.87 mmol) and K<sub>2</sub>CO<sub>3</sub> (654 mg, 4.73 mmol) in DMF (11 mL) under nitrogen was added dropwise a solution of thiophenol (0.15 mL, 1.46 mmol) in DMF (8 mL) over 1 h. The mixture was stirred for an additional 3 h and then concentrated *in vacuo*.

30

The residue was partitioned between  $\text{CHCl}_3$  (50 mL) and  $\text{H}_2\text{O}$  (25 mL). The aqueous phase was separated and extracted with  $\text{CHCl}_3$  ( $3 \times 20$  mL) and the combined organic extracts were dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo*. The residue was purified by column chromatography on basic alumina ( $\text{CHCl}_3$  then 3%  $\text{MeOH}/\text{CHCl}_3$ ) to give the title compound as a yellow oil (178 mg, 23%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.31 (t, 6H,  $J = 7.0$  Hz), 1.40-1.67 (m, 10H), 2.65 (m, 2H), 2.78 (m, 2H), 3.12 (m, 2H), 3.26-3.37 (m, 4H), 3.48 (m, 2H), 3.97-4.09 (m, 4H), 7.61 (m, 1H), 7.68 (m, 2H), 8.06 (m, 1H).

N-[1-[4-Diethoxyphosphoryl-7-(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetradecanyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

Using General Procedure F: Reaction of 4-diethoxyphosphoryl-7-(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetradecane (236 mg, 0.453 mmol) and N-[1-methylene-4-(chloromethylene)phenylene]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (238 mg, 0.551 mmol) followed by silica gel column purification of the reaction products (50% then 80% THF/hexane) gave the title compound as a yellow amorphous solid (305 mg, 73%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.27 (t, 6H,  $J = 7.1$  Hz), 1.43 (br s, 8H), 1.63 (br s, 2H), 2.32 (br s, 2H), 2.55 (m, 2H), 3.13-3.41 (m, 8H), 3.49 (s, 2H), 3.85-4.02 (m, 4H), 4.57 (s, 2H), 4.58 (s, 2H), 7.07-7.22 (m, 6H), 7.51-7.71 (m, 7H), 7.98 (d, 1H,  $J = 7.4$  Hz), 8.04 (m, 1H), 8.41 (d, 1H,  $J = 4.0$  Hz).

N-[1-[7-(2-Nitrobenzenesulfonyl)-1,4,7-triazacyclotetra-decanyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

Using General Procedure E: Reaction of N-[1-[4-diethoxyphosphoryl-7-(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetra-decanyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (300 mg, 0.328 mmol) gave the title compound as a yellow amorphous solid (214 mg, 84%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.34-1.44 (m, 8H), 1.69 (br s, 2H), 2.34 (m, 2H), 2.52 (m, 2H), 2.62 (m, 2H), 2.82 (m, 2H), 3.42 (m, 6H), 4.58 (s, 2H), 4.59 (s, 2H), 7.08-7.24 (m, 6H), 7.52-7.71 (m, 7H), 8.01 (m, 2H), 8.42 (d, 1H,  $J = 4.1$  Hz).

N-[1-(1,4,7-Triazacyclotetra-decanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine Pentahydrobromide Dihydrate

A mixture of N-[1-[7-(2-nitrobenzenesulfonyl)-1,4,7-triazacyclotetra-decanyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (209 mg, 0.268 mmol),  $\text{K}_2\text{CO}_3$  (298 mg, 2.16 mmol), and thiophenol (0.17 mL, 1.7 mmol) in acetonitrile (3 mL) were heated to  $50^\circ\text{C}$  for 16.5 h. The reaction mixture was diluted with  $\text{CH}_2\text{Cl}_2$  (10 mL)

and washed with brine (10 mL). The separated aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 5 mL) and the combined organic extracts were dried ( $\text{MgSO}_4$ ) and evaporated. The residue was purified by column chromatography on basic alumina ( $\text{CHCl}_3$ , then 10%  $\text{MeOH}/\text{CHCl}_3$ ) to give the free base of title compound as a yellow oil (92 mg, 84%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.21-1.62 (m, 10H), 2.40-2.49 (m, 4H), 2.60 (m, 6H), 2.79 (m, 2H), 3.49 (s, 2H), 3.80 (s, 2H), 3.91 (s, 2H), 7.14 (m, 1H), 7.28 (m, 5H), 7.62 (m, 1H), 8.54 (d, 1H,  $J = 4.4$  Hz).

Conversion of the free base (86 mg, 0.21 mmol) to the hydrobromide salt using a saturated solution of  $\text{HBr(g)}$  in  $\text{MeOH}$  (See General Procedure G) followed by drying *in vacuo* at 40 °C for 15.5 h gave the title compound as a white solid (128 mg, 70%):  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ )  $\delta$  1.48 (br s, 6H), 1.82 (m, 4H), 3.22-3.36 (m, 10H), 3.50 (br s, 2H), 4.48 (s, 4H), 4.64 (s, 2H), 7.62 (s, 4H), 7.88 (m, 1H), 7.94 (d, 1H,  $J = 8.0$  Hz), 8.38 (m, 1H), 8.77 (d, 1H,  $J = 5.2$  Hz); FAB-MS  $m/z$  492 ( $\text{MH} + \text{H}^{81}\text{Br}$ ), 490 ( $\text{M} + \text{H}^{79}\text{Br}$ ), 410 ( $\text{M} + \text{H}$ ). Anal. Calcd for  $\text{C}_{25}\text{H}_{39}\text{N}_3 \cdot 5\text{HBr} \cdot 2.5\text{H}_2\text{O} \cdot 0.1\text{Et}_2\text{O}$ : C, 35.20; H, 5.82; N, 8.08; Br, 46.10. Found: C, 35.48; H, 5.66; N, 8.10; Br, 46.05.

#### Example 17

N-[4-[4,7,10,17-Tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7063)

7-Diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene

Using General Procedure H: Reaction of 7-diethoxyphosphoryl-4,10-bis(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene (1.48 g, 2.00 mmol) with thiophenol (with an additional heating of the reaction mixture to 50 °C for 1.5 h after the addition) followed by silica gel column purification of the reaction products (8%  $\text{MeOH}/\text{CHCl}_3$ ) gave the title compound as a light yellow oil (423 mg, 52%):  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.23 (t, 6H,  $J = 7.0$  Hz), 2.50 (br s, 2H), 2.79 (br s, 2H), 3.02-3.15 (m, 10H), 3.82-3.98 (m, 6H), 7.06 (d, 2H,  $J = 7.6$  Hz), 7.54-7.63 (m, 2H), 7.70 (m, 2H), 8.01 (br s, 1H).

N-[4-[7-Diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

Using General Procedure F: Reaction of 7-diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene (410 mg, 0.738 mmol) and N-[1-methylene-4-(chloromethylene)phenylene]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (397 mg, 0.919 mmol) followed by silica gel column purification of the reaction products (50%, 80%, and 90% THF/hexane) gave the title compound as a white amorphous solid (441 mg, 63%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.15 (t, 6H, J = 7.0 Hz), 2.42 (m, 4H), 2.77 (m, 2H), 2.92-3.02 (m, 6H), 3.10 (m, 2H), 3.59 (s, 2H), 3.66-3.91 (m, 6H), 4.58 (s, 2H), 4.59 (s, 2H), 6.94 (d, 1H, J = 7.6 Hz), 7.07-7.14 (m, 6H), 7.22 (d, 1H, J = 7.8 Hz), 7.51-7.72 (m, 8H), 8.00 (d, 1H, J = 7.8 Hz), 8.04 (m, 1H), 8.42 (d, 1H, J = 4.0 Hz).

10 N-[4-[7-Diethoxyphosphoryl-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-2-(aminomethyl)pyridine

A mixture of N-[4-[7-diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (434 mg, 0.456 mmol), K<sub>2</sub>CO<sub>3</sub> (508 mg, 3.68 mmol), and thiophenol (0.28 mL, 2.7 mmol) were heated to 50 °C in CH<sub>3</sub>CN (3.5 mL) under nitrogen for 15 h. Upon cooling, the reaction mixture was partitioned between CHCl<sub>3</sub> (15 mL) and brine (15 mL) and the aqueous layer was separated and extracted with CHCl<sub>3</sub> (3 × 5 mL). The combined organic extracts were dried (MgSO<sub>4</sub>) and concentrated in vacuo and the residue was purified by column chromatography on basic alumina (CHCl<sub>3</sub> then 10% MeOH/CHCl<sub>3</sub>) to give a yellow oil (218 mg, 82%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.16 (t, 6H, J = 7.1 Hz), 2.35 (m, 2H), 2.55 (m, 2H), 2.75 (m, 2H), 2.82 (m, 2H), 2.96-3.08 (m, 6H), 3.16 (m, 2H), 3.68 (s, 2H), 3.69-3.88 (m, 4H), 3.82 (s, 2H), 3.93 (s, 2H), 6.95 (d, 1H, J = 7.6 Hz), 7.00 (d, 1H, J = 7.5 Hz), 7.15-7.34 (m, 6H), 7.50 (m, 1H), 7.65 (m, 1H), 8.56 (d, 1H, J = 4.7 Hz).

25 N-[4-[4,7,10,17-Tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine Hexahydrobromide Hydrate

To a stirred solution of N-[4-[7-diethoxyphosphoryl-4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-2-(aminomethyl)pyridine (211 mg, 0.36 mmol) in acetic acid (0.6 mL) was added a freshly prepared solution of saturated HBr(g) in acetic acid (6 mL) and the reaction mixture was allowed to stir at room temperature for 4 h. Addition of diethyl ether (10 mL) gave a white precipitate that was allowed to settle to the bottom of the flask and the supernatant solution was

decanted. The solid was washed by decantation with MeOH (4 x 5 mL) and ether (6 x 5 mL) and the remaining traces of ether were removed by evaporation under reduced pressure. The product was dried in vacuo at 40 °C for 17 h, to give the title compound as a pale yellow solid (223 mg, 63%): <sup>1</sup>H NMR (D<sub>2</sub>O) δ 3.14-3.36 (m, 10H), 3.55 (m, 4H), 3.75 (m, 2H), 4.45 (s, 2H), 4.50 (s, 2H), 4.64 (s, 2H), 7.22 (m, 2H), 7.53 (s, 4H), 7.70 (m, 1H), 7.95 (m, 1H), 8.00 (d, 1H, J = 7.9 Hz), 8.46 (m, 1H), 8.79 (d, 1H, J = 3.9 Hz); FAB-MS m/z 527 (MH + H<sup>81</sup>Br), 525 (MH + H<sup>79</sup>Br), 445 (M + H). Anal. Calcd for C<sub>27</sub>H<sub>36</sub>N<sub>6</sub>6HBr1.5H<sub>2</sub>O0.2Et<sub>2</sub>O: C, 34.35; H, 4.87; N, 8.65; Br, 49.33. Found: C, 34.57; H, 5.04; N, 8.68; Br, 49.09.

#### Example 18

N-[4-[4,7,10-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD7058)

7-Diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-triene

Using General Procedure H: Reaction of 7-diethoxyphosphoryl-4,10-bis(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-triene

(1.11 g, 1.5 mmol) followed by silica gel column purification of the reaction products (2:5:20 CH<sub>3</sub>OH-Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub> then 1:5 CH<sub>3</sub>OH-CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound as a pale yellow oil (300 mg, 54%): <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 1.21 (t, 6H, J = 7.1 Hz), 1.78-1.92 (br s, 1H), 2.31-2.38 (br m, 2H), 2.56-2.60 (br m, 2H), 2.81-2.98 (br m, 10H), 3.60-3.64 (br m, 2H), 3.75-3.91 (m, 4H), 7.05-7.12 (m, 2H), 7.24-7.29 (m, 2H), 7.60-7.63 (m, 1H), 7.68-7.71 (m, 2H), 8.02-8.06 (m, 1H).

N-[4-[7-Diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

Using General Procedure F: Reaction of 7-diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-triene (290 mg, 0.52 mmol) with N-[1-methylene-4-(chloromethylene)phenylene]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (271 mg, 0.63 mmol) followed by silica gel column purification of the reaction products (1:12:12 CH<sub>3</sub>OH-Et<sub>2</sub>O-CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound (298 mg, 60%) as a



pale yellow solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.17 (t, 6H,  $J = 7.0$  Hz), 2.29-2.45 (br m, 4H), 2.55-2.65 (br m, 2H), 2.71-2.75 (br s, 4H), 2.85-2.91 (br m, 2H), 2.96-2.98 (br m, 2H), 3.57 (s, 2H), 3.67-3.84 (br m, 6H), 4.57-4.61 (br s, 4H), 7.07-7.28 (br m, 10H), 7.55-7.71 (br m, 7H), 7.99-8.02 (m, 2H), 8.42-8.46 (m, 1H).

5        N-[4-[10-(2-Nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine

Using General Procedure E: Reaction of N-[4-[7-diethoxyphosphoryl-10-(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (290 mg, 0.31 mmol) gave the title compound (240 mg, 95%) as a white solid:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.65-1.79 (br s, 1H, coincide with  $\text{H}_2\text{O}$  peak), 2.15-2.19 (br m, 4H), 2.44-2.48 (br m, 2H), 2.61-2.65 (br m, 2H), 2.67-2.71 (br m, 2H), 3.00-3.04 (br m, 2H), 3.10-3.14 (br m, 2H), 3.56-3.60 (br s, 4H), 4.55 (s, 2H), 4.61 (s, 2H), 6.96 (d, 1H,  $J = 7.8$  Hz), 7.02-7.10 (br m, 6H), 7.22-7.28 (br m, 3H), 7.52-7.72 (br m, 7H), 7.96-7.99 (m, 2H), 8.42-8.46 (m, 1H). This was used without further purification.

15        N-[4-[4,7,10-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine Pentahydrobromide Dihydrate

Using General Procedure G: Reaction of N-[4-[10-(2-nitrobenzenesulfonyl)-4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-N-(2-nitrobenzenesulfonyl)-2-(aminomethyl)pyridine (236 mg, 0.29 mmol) followed by alumina column purification of the reaction products (1:99  $\text{CH}_3\text{OH}-\text{CH}_2\text{Cl}_2$  then 1:10  $\text{CH}_3\text{OH}-\text{CH}_2\text{Cl}_2$ ) gave the free base of the title compound (111 mg, 86%) as a pale yellow oil:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  2.24-2.28 (br s, 3H), 2.43-2.50 (br m, 4H), 2.58-2.62 (br m, 2H), 2.73-2.79 (br m, 8H), 2.95-2.98 (br m, 2H), 3.50 (s, 2H), 3.77 (s, 2H), 3.90 (s, 2H), 6.83-6.87 (br m, 3H), 7.05-7.33 (br m, 7H), 7.63-7.67 (m, 1H), 8.54-8.56 (m, 1H).

Conversion of the free base (104 mg, 0.23 mmol) to the hydrobromide salt using a saturated solution of  $\text{HBr(g)}$  in  $\text{CH}_3\text{OH}$  followed by drying of the product *in vacuo*, gave the title compound (101 mg, 52%) as a white solid:  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ )  $\delta$  2.90-2.94 (br m, 2H), 2.97-3.01 (br m, 2H), 3.12-3.16 (br m, 2H), 3.17-3.21 (br m, 2H), 3.24-3.28 (br m, 4H), 3.47-3.51 (br m, 2H), 3.57-3.61 (br m, 2H), 4.38-4.42 (m, 6H), 7.34-7.40 (m, 2H), 7.46-7.60 (m, 8H), 7.90-

7.94 (m, 1H), 8.58-8.62 (m, 1H); FAB-MS  $m/z$  526 (MH + H<sup>81</sup>Br), 524 (MH + H<sup>79</sup>Br), 444 (M + H, 100); Anal. Calcd. for C<sub>28</sub>H<sub>42</sub>N<sub>3</sub>Br<sub>5</sub>•2.5H<sub>2</sub>O: C, 37.65; H, 5.30; N, 7.84; Br, 44.73. Found: C, 37.53; H, 5.26; N, 7.79; Br, 44.75.

5            Table 2

Compound	Inhibition of mAb 12G5 binding
	IC <sub>50</sub> <sup>a</sup> (ng/ml)
AMD3100	27
AMD3465	3
AMD7049	52
AMD7050	1
AMD7051	7
AMD7058	>1000
AMD7059	>1000
AMD7063	9
SDF-1α <sup>b</sup>	270

<sup>a</sup> Inhibition of mAb 12G5 binding to CXCR-4 in SUP-T1 cells.

<sup>b</sup> Natural ligand for CXCR-4 (Bleul *et al. Nature*, 382: 829-832 (1996); Oberlin *et al., Nature*, 382:833-835 (1996)).

Table 3

Compound	% Inhibition of Ca <sup>2+</sup> flux (conc) or IC <sub>50</sub> <sup>a</sup> (ng/ml)
AMD3100	5
AMD3465	1
AMD7049	100% (1 µg/ml)
AMD7050	100% (1 µg/ml)
AMD7051	100% (1 µg/ml)
AMD7058	44% (1 µg/ml)
AMD7059	36% (1 µg/ml)
AMD7063	100% (1 µg/ml)

<sup>a</sup> Inhibition of Signal transduction (increasing intracellular Ca<sup>2+</sup> flux) induced by SDF-1 $\alpha$  binding to CXCR-4 on SUP-T1 cells.

Each of the following compounds, including AMD 3484 (see Fig.28), were synthesized according to procedures in Bridger *et al.*, *J. Med. Chem.* 1995, 38, 366-378; *J. Med. Chem.* 1996, 39, 109-119 and U.S. Pat. No. 5,583,131, U.S. Pat. No. 5,698,546 and U.S. Pat. No. 5,817,807, which are each incorporated in their entirety by reference herein.

5

Example 191-[2,6-Dimethoxypyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane Tetrahydrobromide  
(AMD 7032)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.78 (m, 2H), 1.88-1.92 (m, 2H), 2.59-3.03 (m, 16H), 3.60 (s, 2H), 3.91 (s, 6H), 6.44 (s, 2H); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 26.75, 27.91, 48.34, 49.21, 49.89, 50.96, 52.01, 52.86, 54.88, 57.15, 57.53, 59.42, 142.65, 157.42, 166.42; FAB MS *m/z* 352 (M+H); Anal. (C<sub>18</sub>H<sub>33</sub>N<sub>5</sub>O<sub>2</sub> 4.1 HBr 0.25 H<sub>2</sub>O); Calc. C, 31.44; H, 5.51; N, 10.18; Br, 47.64. Found C, 31.17; H, 5.61; N, 9.92; Br, 47.54.

10

Example 201-[2-Chloropyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane Tetrahydrochloride  
Monohydrate (AMD 7048)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.92 (m, 2H), 2.12 (m, 2H), 2.77-2.80 (m, 4H), 2.96-3.39 (m, 12H), 3.85 (s, 2H), 7.33 (d, 1H, *J* = 5.4 Hz), 7.44 (s, 1H), 8.40 (d, 1H, *J* = 5.4 Hz); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 24.75, 27.59, 47.40, 47.55, 49.11, 49.23, 52.12, 52.40, 53.81, 54.42, 56.98, 126.97, 128.30, 151.90, 152.34, 153.78; FAB MS *m/z* 326 (M+H); Anal. (C<sub>16</sub>H<sub>28</sub>N<sub>5</sub>Cl<sub>4</sub>·2HCl·0.5HOAc·1.1H<sub>2</sub>O); Calc. C, 38.61; H, 6.94; N, 13.24; Cl, 34.86. Found C, 38.63; H, 6.94; N, 13.52; Cl, 34.61.

20

Example 211-[2,6-Dimethylpyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane Pentahydrobromide  
Dihydrate (AMD 7060)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.77 (m, 2H), 1.93 (m, 2H), 2.48 (s, 6H), 2.61-3.00 (m, 16H), 3.61 (s, 2H), 7.07 (s, 2H); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 25.30, 26.22, 27.49, 47.75, 48.65, 49.43, 50.41, 51.58, 52.19, 54.09, 56.63, 58.46; FAB MS *m/z* 320 (M+H); Anal. (C<sub>18</sub>H<sub>33</sub>N<sub>5</sub>·5HBr·0.5HOAc·1.7H<sub>2</sub>O); Calc. C, 29.08; H, 5.57; N, 8.92; Br, 50.91. Found C, 29.04; H, 5.60; N, 8.73; Br, 50.87.

30

Example 221-[2-Methylpyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane Pentahydrobromide  
Dihydrate (AMD 7061)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 2.01-2.08 (m, 2H), 2.22 (m, 2H), 2.70-2.72 (m, 2H), 2.77 (s, 3H), 2.91-2.92 (m, 2H), 3.33-3.52 (m, 12H), 4.00 (s, 2H), 7.86-7.89 (m, 2H), 8.56 (d, 1H, *J* = 5.7 Hz); FAB MS *m/z* 306 (M+H); Anal. (C<sub>17</sub>H<sub>31</sub>N<sub>5</sub>·4.9HBr·0.3HOAc·2.1 H<sub>2</sub>O); Calc. C, 27.9; H, 5.49; N, 9.24; Br, 51.67. Found C, 28.08; H, 5.50; N, 9.56; Br, 51.56.

40

Example 231-[2,6-Dichloropyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane Trihydrochloride Bishydrate (AMD 3451)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.83-1.88 (m, 2H), 2.04-2.08 (m, 2H), 2.58-2.62 (m, 2H), 2.79-2.81 (m, 2H), 3.12-3.44 (m, 12H), 3.69 (s, 2H), 7.30 (s, 2H); <sup>13</sup>C NMR (D<sub>2</sub>O) □□36.26, 37.69, 55.26, 56.18, 58.33, 58.56, 58.92, 59.23, 63.57, 65.44, 70.72, 140.37, 166.14, 167.37; FAB MS *m/z* 360 (M+H). Anal. (C<sub>16</sub>H<sub>34</sub>N<sub>5</sub>Cl<sub>3</sub>O<sub>2</sub>): Calc. C, 38.00; H, 6.78; N, 13.85; Cl, 35.05. Found: C, 38.33; H, 6.42; N, 13.88; Cl, 35.43.

Example 241-[2-Chloropyrid-5-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane Tetrahydrochloride Hemihydrate (AMD 3454)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.96-2.09 (br m, 4H), 3.02-3.17 (m, 4H), 3.19-3.28 (br m, 8H), 3.40 (s, 4H), 4.10 (s, 2H), 7.40 (d, 1H, *J* = 8.2 Hz), 7.80 (d, 1H, *J* = 8.2 Hz), 8.27 (s, 1H); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 19.36, 19.47, 38.17, 38.64, 39.06, 41.74, 41.88, 42.18, 45.66, 48.29, 54.62, 125.59, 126.69, 142.79, 150.77, 151.75; FAB-MS *m/z* 326 (M + H). Anal. Calcd for C<sub>16</sub>H<sub>28</sub>N<sub>5</sub>Cl<sub>4</sub>HCl<sub>0.5</sub>H<sub>2</sub>O: C, 39.98; H, 6.92; N, 14.57; Cl, 36.87. Found: C, 40.36; H, 7.06; N, 14.56; Cl, 36.92.

General Procedures A, B and C were used to prepare the following compounds:

Example 25N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-purine Pentahydrobromide DiHydrate (AMD3472)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.88-2.05 (br m, 4H), 3.06-3.22 (br m, 8H), 3.27-3.44 (br m, 8H), 4.22 (s, 2H), 5.59 (s, 2H), 7.29 (s, 4H), 8.80 (s, 1H), 9.11 (s, 1H), 9.28 (s, 1H); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 18.39, 19.25, 37.24, 37.55, 37.71, 41.13, 41.37, 41.71, 44.41, 47.73, 54.87, 129.45, 131.81, 132.53, 136.67, 140.96, 147.88, 152.46, 154.37; FAB-MS *m/z* 423 (M + H). Anal. Calcd for C<sub>23</sub>H<sub>34</sub>N<sub>8</sub>.5HBr.2H<sub>2</sub>O.0.5CH<sub>3</sub>CO<sub>2</sub>H: C, 32.27; H, 5.07; N, 12.54; Br, 44.73. Found: C, 32.66; H, 4.81; N, 12.41; Br, 44.58.

Example 261-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-phenylpiperazine Pentahydrobromide Hydrate (AMD3526)

<sup>1</sup>H NMR (D<sub>2</sub>O) δ 1.88-2.06 (br m, 4H), 3.11-3.53 (br m, 24H), 4.30 (s, 2H), 4.32 (s, 2H), 6.89-6.97 (m, 3H), 7.19-7.24 (m, 2H), 7.49 (s, 4H); <sup>13</sup>C NMR (D<sub>2</sub>O) δ 18.74, 19.37, 37.34, 41.47, 41.76, 42.03, 44.31, 47.45, 48.26, 51.16, 58.48, 59.29, 118.18, 122.34, 129.99, 130.37, 131.53, 131.85, 132.62, 148.47; FAB-MS m/z 547 (M + H<sup>81</sup>Br), 545 (M + H<sup>79</sup>Br), 465 (M + H). Anal. Calcd for C<sub>28</sub>H<sub>44</sub>N<sub>6</sub>.5HBr.H<sub>2</sub>O: C, 37.90; H, 5.79; N, 9.47; Br, 45.03. Found: C, 37.72; H, 5.98; N, 9.38; Br, 46.78.

Table 4

Compound	IC <sub>50</sub> <sup>a</sup> (μg/mL)
AMD3451	8.9
AMD3472	45.4
AMD3454	32.3
AMD3526	82
AMD3100	>100

<sup>a</sup> 50% Inhibitory Concentration (IC<sub>50</sub>)(μg/mL) exhibited by AMD compounds against infection of U87.CD4.CCR5 by HIV-1 BaL

### Example 27

#### Inhibition of Collagen-Induced Arthritis

A compound as used in the present invention demonstrated inhibition of collagen-induced arthritis (CIA) in a mutant mouse model.

### METHODOLOGY

#### Experimental Animal Treatment

The control group consisted of ten mice that were injected with collagen as discussed below. The treatment group consisted of eight mice which were also injected with collagen and were further treated by administering a compound of (1,1'-[1,4-phenylenebis(methylene)])bis-1,4,8,11-tetraazacyclotetradecane) (AMD 3100; see figure 27) intravenously using osmotic pumps (200μl, Alza, 0.5 μl/hr) at a concentration of 5 mg/ml over a 14-day period following collagen injection.

25 Mutant Mice

The generation and the basic characteristics of the mutant mouse strain (129/Sv/Ev) with a disruption in the gene coding for the  $\alpha$ -chain of the IFN- $\gamma$  receptor (IFN- $\gamma$  RKO) have been described (Huang S. *et al.*, *Science* 259:1742, (1993)). These IFN- $\gamma$  RKO mice were back-crossed with DBA/1 wild-type mice for 10 generations to obtain the DBA/1 IFN- $\gamma$  RKO mice used in the present study. IFN- $\gamma$  RKO and wild-type mice were bred in the Experimental Animal Centre of the University of Leuven. The experiments were performed in 8- to 12-week old mice, but in each experiment, the mutant and wild-type mice were age-matched with a 5 day limit. The male to female ratio was kept between 0.8 and 1.3 in each experimental group.

#### Induction of Collagen-Induced Arthritis and Clinical Assessment of Arthritis

Collagen-induced arthritis was carried out in the following manner (see: Vermeire *et al.*, *Int. J. Immunol.* 158:5507-5513, (1997)). Native chicken collagen type II (EPC, Owensville, MO) was dissolved in 0.05 M acetic acid at 2 mg/ml by stirring overnight at 6°C, and emulsified in an equal volume of incomplete (IFA) or complete Freund's adjuvant (CFA) containing 1.5 mg/ml heat killed *Mycobacterium butyricum* (Difco, Detroit, MI). Mice were sensitized with a single 100  $\mu$ l intradermal injection of the emulsion at the base of the tail. Mice were examined daily for signs of arthritis. The disease severity was recorded following a scoring system for each limb. Score 0: normal; score 1: redness and /or swelling in one joint; score 2: redness and/or swelling in more than one joint; score 3: redness and/or swelling in the entire paw; score 4: deformity and/or ankylosis.

#### Histological Examination

Spleens and fore and hind limbs were fixed in buffered saline - B5fixative (10% formalin with quicksilver). Alternatively, tissues were fixed in 10% formalin or pure methanol (see: Vermeire *et al.*, *J. Immunol.*, 158, 5507-5513, 1997). Limbs were subsequently decalcified overnight with formic acid. Four-micron thick paraffin sections were stained with hematoxylin and eosin. Severity of arthritis was evaluated using three parameters: infiltration of mono- and polymorphonuclear cells, hyperplasia of the synovium and pannus formation. Each parameter was scored on a scale from 0 to 3 (absent; weak, moderate and severe).

#### In Vivo Antibody Treatments

Monoclonal antibodies were produced from hybridomas grown by intraperitoneal inoculation in Pristane-primed athymic nude mice (nu/nu of NMRI background). Neutralizing monoclonal antibody against MuIFN- $\gamma$  (F3, rate IgG<sub>2a</sub>) was purified by affinity chromatography

on a mouse anti-rat  $\kappa$  chain monoclonal antibody (Billiau A. *et al.*, *J. Immunol.* 140:1506, (1988)). The neutralizing titer (end-point dilution corresponding to 50% neutralization of the antiviral effect of 30 units/ml of mouse IFN- $\gamma$  on mouse I929 cells challenged with mengovirus) was  $10^{53}$  U/ml (IgG content, 1.4 mg/ml). A neutralizing rate IgG<sub>24</sub> antibody against murine IL-12 was produced using hybridoma C17.8 (kindly provided by Dr. G. Trinchieri, Wistar Institute, Philadelphia, PA). The antibody was purified by affinity chromatography on protein G (Pharmacia, Uppsala, Sweden). Antibody against murine IL-6 was prepared from ascites fluid from thymus-less nude mice inoculated with the 20F3 (rat x mouse) hybridoma (American Type Culture Collection, Rockville, MD). This rat IgG antibody was purified by affinity chromatography on an anti-rat  $\kappa$  chain monoclonal antibody-Sepharose column. The neutralizing titer (endpoint dilution corresponding to 50% neutralization of the cell growth effect of 10 U of murine IL-6 per ml) was  $10^{55}$  (IgG content: 2.9 mg/ml).. Irrelevant rat IgG<sub>24</sub> was used as an isotope control and was prepared from ascites fluid of a rat plasmocytoma (obtained through the courtesy of Dr. H. Bazin, University of Louvain, Medical School, Brussels, Belgium). The IgG was purified by anion exchange chromatography on Hiload Q Sepharose and gel filtration on Superdex 200 (Pharmacia). Batches of anti-IFN- $\gamma$ , anti-IL-12, anti-IL-6 and irrelevant IgG<sub>24</sub> were tested for endotoxin content by a chromogenic *Limulus* amoebocyte lysate assay (KabiVitrum, Stockholm, Sweden) and were found to contain less than 2 ng/ml endotoxin. Injections were given in 200  $\mu$ l of pyrogen-free saline.

## RESULTS

Following 14 days after treatment, 7 of the ten mice in the control group demonstrated arthritis, while only 1 of the 8 treated animals demonstrated disease. See Figure 29. The single treated animal did not develop arthritic pathology until after 20 days post-treatment. Additionally, the treated animals compared with the control animals did not demonstrate any significant body weight loss. See Figure 30. Further, the treated animals maintained body weight consist with healthy animals not injected with collagen (curve not shown).

### Example 28

#### Treatment of Glioblastoma

Compounds of the present invention, such as AMD 3100 may be used in the treatment of glioblastomas, fibromas, astrocytomas or myelomas affecting the central nervous system.



The compounds may be used according to standard clinical practice and procedures, using dosages as provided in the foregoing examples and according to clinical end points, such as imaging, immunological and other methodologies.

For example, the etiology or association of chemokine receptor binding in the proliferation of glioblastoma tumor cells has been reported by Sehgal *et al.*, *J. of Surg. Oncol.* 69:99-104 (1998) ("Sehgal I") and Sehgal *et al.*, *J. of Surg. Oncol.* 69:239-248 (1998) ("Sehgal II"). The role of CXCR4 of its binding to its receptor appears to play a significant role in the formation and/or proliferation of glioblastoma cells. The inhibition of the binding by CXCR4 to its natural receptor ligand by compounds of the present invention, such as AMD 3100, offer a new drug in the treatment tumors of central nervous system that are mediated or associated with chemokines, such as CXCR4.

#### Example 29

##### Treatment of Non-Small Cell Lung Cancer

Compounds of the present invention, such as AMD 3100, may be used in the treatment of non-small cell lung cancer. The compounds may be used according to standard clinical practice and procedures, using dosages as provided in the foregoing examples and according to clinical end points, such as imaging, immunological and other methodologies.

For example, CXC chemokines have been found to regulate or are associated with the regulation of angiogenesis in non-small cell lung cancer (see: Arenberg, *et al.*, *J. of Leukocyte Biol.*; 62:554-562 (1997); and Moore *et al.* *TCM*, vol 8(2): 51-58 (1998) Elsevier Science, Inc.). The role of CXC chemokines and the binding to their respective receptors appear to play a significant role in the formation and/or proliferation of non-small cell lung cancer promoted by an increase in angiogenic activity. The inhibition of the binding of these CXC chemokines to their natural receptor ligands by compounds of the present invention, such as AMD 3100, offer a new drug in the treatment tumors such as non-small cell lung cancer that are mediated or associated with increased levels of chemokines.

#### Example 30

N-[4-(1,7-Diazacyclotetradecanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine

The compounds of the present invention further include compounds of formula I, where V has 2 to 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other. An example of such compound, includes:

N-[4-(1,7-Diazacyclotetradecanyl)-1,4-phenylenebis(methylene)]-2-  
5 (aminomethyl)pyridine (AMD-Exp 1; see Figure 34).

AMD-Exp 1, Exp-2 and Exp3 may each be prepared by modification of our previously published strategies (Bridger *et al. J. Med. Chem.* 1995, 38, 366) for preparing carbon-linked azamacrocycles and is briefly summarized as follows: The aforementioned intermediate, N-[1-methylene-4-(chloromethylene)phenylene]-N-(2-nitrobenzenesulfonyl)-2-aminomethyl)pyridine  
10 (or the Dep -protected intermediate) is reacted with the sodium salt of diethylmalonate to give the corresponding diester.

Reduction of the diester to the corresponding diol, followed by derivatization with methanesulfonyl chloride gave the dimesylate. Nucleophilic displacement of the dimesylate with sodium cyanide gives the requisite dinitrile which was reduced to the diamine with  
15 borane.THF and derivatized with 2-nitrobenzenesulfonyl chloride for the impending macrocyclization reaction.

Macrocyclization with an appropriately derivatized diol (such as 1,7-heptanediol di-p-tosylate) and subsequent deprotection as described in previous examples gave, for example, AMD-Exp 1. Similarly, 1,3-phenylenediethanol and 2,6-pyridine diethanol would give AMD-  
20 Exp 2 and AMD-Exp-3 respectively.

Further, AMD-Exp 1, AMD-Exp 2, and AMD-Exp 3 may each be prepared into a pharmaceutical compositions comprising a therapeutically effective amount of the compound in a pharmaceutically acceptable carrier.

Additionally, AMD-Exp 1, AMD-Exp 2, and AMD-Exp 3 may each be used according  
25 to the foregoing methods for the treatment of a number of chemokine-mediated diseases and conditions, including: infection with HIV or FIV; a disease by the regulation of endothelial cell function; a disease relating to vascularization of the gastrointestinal tract; haematopoiesis; or cerebellar development; a disease relating to basal leukocyte trafficking or the extravasation and tissue infiltration of leukocytes in response to inciting antigens; compound effectively binds to a  
30 chemokine receptor; of inflammatory disease; cancer; central nervous system developmental disease; HIV; FIV; vasculature development disease; cardiogenesis developmental disease;

haematopoiesis and other chemokine mediated diseases or disorders.

Additional examples of such specific diseases or conditions that may be treated using AMD-Exp 1, further include: inflammation due to arthritis or multiple sclerosis; cancers associated with: solid tumors; lymphoma; metastatic tumors; glioblastoma tumors; and other carcinoma tumors; non-small cell lung cancer; lung cancer; breast cancer; prostate cancer; and cancer of other organs. Further, disorders or conditions that may be treated using AMD-Exp 1, includes: disorders treated by inhibiting or promoting angiogenesis or by inducing stasis of angiogenesis.

10

### Example 31

N-[7-(4,10-Diazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine

The compounds of the present invention further include compounds of formula I, where V has 2 to 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other. An example of such compound, includes:

N-[7-(4,10-Diazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD-Exp 2; see Figure 36).

AMD-Exp 2 and AMD-Exp 3 may each be prepared according to Example 30 above. Further, AMD-Exp 2 may be prepared into a pharmaceutical composition comprising a therapeutically effective amount of the compound in a pharmaceutically acceptable carrier.

Additionally, AMD-Exp 2 may be used according to the foregoing methods for the treatment of a number of chemokine-mediated diseases and conditions, including: infection with HIV or FIV; a disease by the regulation of endothelial cell function; a disease relating to vascularization of the gastrointestinal tract; haematopoiesis; or cerebellar development; a disease relating to basal leukocyte trafficking or the extravasation and tissue infiltration of leukocytes in response to inciting antigens; compound effectively binds to a chemokine receptor; of inflammatory disease; cancer; central nervous system developmental disease; HIV; FIV; vasculature development disease; cardiogenesis developmental disease; haematopoiesis and other chemokine mediated diseases or disorders.

30

Additional examples of such specific diseases or conditions that may be treated using AMD-Exp 2, further include: inflammation due to arthritis or multiple sclerosis;

cancers associated with: solid tumors; lymphoma; metastatic tumors; glioblastoma tumors; and other carcinoma tumors; non-small cell lung cancer; lung cancer; breast cancer; prostate cancer; and cancer of other organs. Further, disorders or conditions that may be treated using AMD-Exp 2, includes: disorders treated by inhibiting or promoting angiogenesis or by  
5 inducing stasis of angiogenesis.

The active compounds may be administered in the form of a pharmaceutical composition formulated according to well known principles and incorporating the compound, preferably in unit dose form, in combination with a pharmaceutically acceptable diluent, carrier  
10 or excipient. Such compositions may be in the form of solutions or suspensions for injection, or irrigation or be in capsule, tablet, dragee, or other solid composition or as a solution or suspension for oral administration or formulated into pessaries or suppositories or sustained release forms of any of the above for implantation. Suitable diluents, carriers, excipients and other components are well known. It may be desirable also to formulate a composition for  
15 topical administration such as an ointment or cream. The compounds of the invention may be used, in the form of a composition or alone.

The pharmaceutical compositions according to the invention may be formulated in unit dosages determined in accordance with conventional pharmacological methods, suitably to provide active compounds in the dosage range in humans or animals of from 0.01 to 100 mg/kg  
20 body weight per day, in a single dose or in a number of smaller doses. Preferred dosage ranges are 0.01 to 30 mg/kg body weight per day intravenous (iv) or intraperitoneal (ip). Other active compounds may be used in the compositions or such active compounds or supplemental therapy may be included in a course of treatment. The pharmaceutical compositions are useful for treatment of a patient comprising an effective therapeutic amount of the novel compound,  
25 where said compound effectively binds to a chemokine receptor.

The present invention further contemplates the use of these compositions in the manufacture of a medicament for the treatment of HIV-or FIV-infected patients and/or the treatment of a disease by the regulation of endothelial cell function and/or the treatment of a disease relating to vascularization of the gastrointestinal tract; haematopoiesis; or cerebellar  
30 development.

In a method for treating a patient infected with HIV or FIV, the pharmaceutical

composition is administered to said patient as a therapeutically effective amount of a pharmaceutical composition in a pharmaceutically acceptable carrier. In a method of treating a patient with a disease related to the regulation of endothelial cell function, the pharmaceutical composition is administered to said patient as a therapeutically effective amount of a pharmaceutical composition in a pharmaceutically acceptable carrier. The present invention further contemplates methods of treating a patient with a disease relating to vascularization of the gastrointestinal tract; haematopoiesis; or cerebellar development, by administering to said patient a therapeutically effective amount of a pharmaceutical composition in a pharmaceutically acceptable carrier.

10       The present invention further contemplates a method of treating a patient with a disease relating to basal leukocyte trafficking or the extravasation and tissue infiltration of leukocytes in response to inciting antigens, by administering to said patient a therapeutically effective amount of a pharmaceutical composition in a pharmaceutically acceptable carrier. The present method also contemplates treating a patient, by administering to said patient a therapeutically effective amount of a pharmaceutical composition in a pharmaceutically acceptable carrier, wherein said compound effectively binds to a chemokine receptor.

15       The present invention further contemplates pharmaceutical compositions and methods of use for the treatment of humans or animals for: renal allograft rejection; inflammatory disease; cancer; central nervous system developmental disease; HIV; vasculature development disease; haematopoiesis and other chemokine mediated diseases or disorders. This invention further provides for the treatment of diseases, which include, but are not limited to: arthritis; multiple sclerosis; dementia from HIV or FIV infection, Parkinson's disease, Alzheimer's disease and inflammatory diseases. The pharmaceutical compositions and methods of use of the present invention further provide for the treatment of cancers, that include, but are not limited to those associated with: solid tumors; lymphoma; metastatic tumors; glioblastoma tumors; and other carcinomas tumors. The pharmaceutical compositions of the present invention are useful for the treatment of cancers that include, but are not limited to: non-small cell lung cancer; lung cancer; breast cancer; prostate cancer; and cancer of other organs.

20       Other diseases or disorders that are contemplated to be treated with the pharmaceutical compositions of the present invention, include, but are not limited to: disorders treated by


inhibiting or promoting angiogenesis or by inducing stasis of angiogenesis; developmental disorders mediated by chemokines.

The present invention further provides methods for the prevention of a disease or disorder in a patient by administering a therapeutically effective dosage of the pharmaceutical  
5 compositions of the present invention to a patient over a period of time sufficient to effectively prevent the disease or disorder.

CLAIMS

1. A macrocyclic compound of general formula I,

5



wherein V is a cyclic polyamine moiety having a total of 9 to 24 members and from 2 to 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic ring;

R<sup>1</sup> to R<sup>7</sup> may be the same or different and are independently selected from hydrogen or straight, branched or cyclic C<sub>1-6</sub> alkyl;

R<sup>8</sup> is a heterocyclic group, a substituted aromatic group, or a mercaptan group;

Ar is an aromatic or heteroaromatic ring each optionally substituted at single or multiple positions with electron-donating or withdrawing groups;

x is 1 or 2;

and the acid addition salts and metal complexes thereof.

2. The macrocyclic compound of claim 1, wherein V is a 14 to 20 membered fused or unfused ring system.

3. A compound of claim 2, which is:


N-[4-(1,7-Diazacyclotetradecanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD-Exp 1).

4. A compound of claim 2, which is:

N-[7-(4,10-Diazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD-Exp 2).

30

5. A macrocyclic compound of general formula I,



wherein V is a cyclic polyamine moiety having a total of 9 to 24 members and from 3 to  
 5 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon  
 atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic  
 ring;

R<sup>1</sup> to R<sup>7</sup> may be the same or different and are independently selected from hydrogen or  
 straight, branched or cyclic C<sub>1-6</sub> alkyl;

10 R<sup>8</sup> is a heterocyclic group, a substituted aromatic group, or a mercaptan group;

Ar is an aromatic or heteroaromatic ring each optionally substituted at single or multiple  
 positions with electrons-donating or withdrawing groups;

x is 1 or 2;

and the acid addition salts and metal complexes thereof.

15

6. The macrocyclic compound of claim 5, wherein V is a 14 to 20 membered fused  
 or unfused ring system.

7. The macrocyclic compound of claim 5, wherein V is a 4,7,10,17-  
 20 tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl system or a 1,4,7-triazacyclotetra-decanyl  
 or a 4,7,10-triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl system, or a 1,4,8,11-  
 tetraazacyclotetradecanyl system or a derivative thereof.

8. A compound of claim 5, which is:

25 N-[4-(1,4,7-Triazacyclotetra-decanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine  
 (AMD 7049);

N-[7-(4,7,10,17-Tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-  
 phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7050);

30 N-[7-(4,7,10-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-  
 phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7051);




- N-[4-[4,7,10-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7058);
- N-[1-(1,4,7-Triazacyclotetra-decanyl)-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7059);
- 5 N-[4-[4,7,10,17-Tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl]-1,4-phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 7063);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-methyl)pyridine (AMD 3465);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-*N*-methyl-2-
- 10 (aminomethyl)pyridine (AMD 3538);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-amino-methylpyridine (AMD 3500);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-3-(amino-methyl)pyridine (AMD 3499);
- 15 N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-(2-amino-methyl-5-methyl)pyrazine (AMD 3498);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-ethyl)pyridine (AMD 3497);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-
- 20 methyl)thiophene (AMD 3516);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-(amino-ethyl)mercaptan (AMD 3530);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-2-amino-benzylamine (AMD 3517);
- 25 N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-amino-benzylamine (AMD 3544);
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-(amino-ethyl)imidazole (AMD 3543); or
- N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-benzylamine (AMD
- 30 3529);
- N-[3-(3,6,17-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-

phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD 8630);  
N-[3-(3,6,17-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,3-  
phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD8631);  
N-[4-(4,7,17-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-  
5 phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD7450);  
N-[7-(4,7,17-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-  
phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD7463);  
N-[6-(3,6,9-Triazabicyclo[11.3.1]pentadeca-1(15),11,13-trienyl)-1,3-  
phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD7097); or  
10 N-[7-(4,10,17-Triazabicyclo[13.3.1]heptadeca-1(17),13,15-trienyl)-1,4-  
phenylenebis(methylene)]-2-(aminomethyl)pyridine (AMD-Exp 3); or  
the acid addition salts and metal complexes thereof.

9. A macrocyclic compound of general formula V,

15



wherein  $V^2$  is a cyclic polyamine moiety having a total of 9 to 24 members and from 3  
to 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon  
atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic  
20 ring;

$R^9$  and  $R^{10}$  may be the same or different and are independently selected from hydrogen  
or straight, branched or cyclic  $C_{1-6}$  alkyl;

$Ar^2$  is an aromatic or heterocyclic ring each optionally substituted at single or multiple  
positions with electron-donating or withdrawing groups and/or aromatic and heterocyclic  
25 groups and their alkyl derivatives thereof; and the acid addition salts and metal complexes.

10. The compound of claim 9, wherein  $V^2$  is a cyclam or substituted cyclam system  
or a 1,4,8,11-tetraazacyclotetradecanyl system or a 4,7,10,17-tetraazabicyclo[13.3.1]heptadeca-  
1(17),13,15-trienyl system.

30

11. The compound of claim 10, wherein Ar<sup>2</sup> is a heterocyclic or substituted heterocyclic.

12. The compound of claim 11, wherein the or heterocyclic substituents are  
5 independently selected from the group consisting of: alkoxy; alkyl; halogen and hydrogen.

13. The compound of claim 12, which is:  
1-[2,6-Dimethoxypyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane (AMD 7032);  
1-[2-Chloropyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane (AMD 7048);  
10 1-[2,6-Dimethylpyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane (AMD 7060);  
1-[2-Methylpyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane (AMD 7061);  
1-[2,6-Dichloropyrid-4-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane (AMD 3451); or  
1-[2-Chloropyrid-5-yl(methylene)]-1,4,8,11-tetraazacyclotetradecane (AMD 3454); or  
the acid addition salts and metal complexes thereof.

15

14. The compound of claim 10, wherein Ar<sup>2</sup> is an aromatic or substituted aromatic ring.

15. The compound of claim 14, wherein the aromatic substituents are independently  
20 selected from the group consisting of: alkoxy; alkyl; halogen; hydrogen; aromatic; heterocyclic or substituted heterocyclic.

16. The compound of claim 15, which is:  
N-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-purine (AMD 3472);  
25 1-[1,4,8,11-Tetraazacyclotetradecanyl-1,4-phenylenebis(methylene)]-4-phenylpiperazine  
(AMD3526); or  
7-[4-methylphenyl (methylene)]-4,7,10,17-Tetraazabicyclo[13.3.1]heptadeca-1(17),13,15-triene  
(AMD 3484).


30 17. A pharmaceutical composition comprising an effective therapeutic amount of the compound of any one of claims 1 to 16.

18. A composition according to claim 17, in unit dosage form.
19. The use of a composition of claim 17, in the manufacture of a medicament for  
5 the treatment of HIV- or FIV-infected patients.
20. The use of a compound of claim 17, in the manufacture of a medicament for the treatment of a disease by the regulation of endothelial cell function.
- 10 21. The use of a composition of claim 17, in the manufacture of a medicament for the treatment of a disease relating to vascularization of the gastrointestinal tract; haematopoiesis; or cerebellar development.
- 15 22. A method of treating a patient infected with HIV or FIV, by administering to said patient a therapeutically effective amount of a pharmaceutical composition comprising the composition of claim 17 in a pharmaceutically acceptable carrier.
- 20 23. A method of treating a patient with a disease related to the regulation of endothelial cell function, by administering to said patient a therapeutically effective amount of a pharmaceutical composition comprising the composition of claim 13 in a pharmaceutically acceptable carrier.
- 25 24. A method of treating a patient with a disease relating to vascularization of the gastrointestinal tract; haematopoiesis; or cerebellar development, by administering to said patient a therapeutically effective amount of a pharmaceutical composition comprising the composition of claim 17 in a pharmaceutically acceptable carrier.
- 30 25. A method of treating a patient with a disease relating to basal leukocyte trafficking or the extravasation and tissue infiltration of leukocytes in response to inciting antigens, by administering to said patient a therapeutically effective amount of a pharmaceutical composition comprising the composition of claim 17 in a pharmaceutically acceptable carrier.

26. A pharmaceutical composition useful for treatment of a patient comprising an effective therapeutic amount of the compound of any one of claims 1 to 16, wherein said compound effectively binds to a chemokine receptor.

27. A method of treating a patient, by administering to said patient a therapeutically effective amount of a pharmaceutical composition comprising the composition of claim 17 in a pharmaceutically acceptable carrier, wherein said compound effectively binds to a chemokine receptor.

28. A pharmaceutical composition comprising a therapeutically effective amount of a compound of general formula I,



wherein V is a cyclic polyamine moiety having a total of 9 to 24 members and from 2 to 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic ring;

R<sup>1</sup> to R<sup>7</sup> may be the same or different and are independently selected from hydrogen or straight, branched or cyclic C<sub>1-6</sub> alkyl;

R<sup>8</sup> is a heterocyclic group, a substituted aromatic group, or a mercaptan group;


Ar is an aromatic or heteroaromatic ring each optionally substituted at single or multiple positions with electrons-donating or withdrawing groups;

x is 1 or 2;

and the acid addition salts and metal complexes thereof, and

wherein said pharmaceutical composition is useful for the treatment of inflammatory disease; cancer; central nervous system developmental disease; HIV; FIV; vasculature development disease; haematopoiesis and other chemokine mediated diseases or disorders.

29. A pharmaceutical composition comprising a therapeutically effective amount of a compound of general formula I,



5

wherein V is a cyclic polyamine moiety having a total of 9 to 24 members and from 3 to 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic ring;

10  $R^1$  to  $R^7$  may be the same or different and are independently selected from hydrogen or straight, branched or cyclic  $C_{1-6}$  alkyl;

$R^8$  is a heterocyclic group, a substituted aromatic group, or a mercaptan group;

Ar is an aromatic or heteroaromatic ring each optionally substituted at single or multiple positions with electrons-donating or withdrawing groups;


15 x is 1 or 2;

and the acid addition salts and metal complexes thereof, and

wherein said pharmaceutical composition is useful for the treatment of inflammatory disease; cancer; central nervous system developmental disease; HIV; FIV; vasculature development disease; haematopoiesis and other chemokine mediated diseases or disorders.

20

30. A pharmaceutical composition comprising a therapeutically effective amount of a compound of general formula V,



25

wherein  $V^2$  is a cyclic polyamine moiety having a total of 9 to 24 members and from 3 to 6 optionally substituted amine nitrogens spaced by two or more optionally substituted carbon atoms from each other, and which may optionally comprise a fused aromatic or heteroaromatic ring;

30  $R^9$  and  $R^{10}$  may be the same or different and are independently selected from hydrogen or straight, branched or cyclic  $C_{1-6}$  alkyl;

Ar<sup>2</sup> is an aromatic or heterocyclic ring each optionally substituted at single or multiple positions with electron-donating or withdrawing groups and/or aromatic and heterocyclic groups and their alkyl derivatives thereof; and the acid addition salts and metal complexes, and

wherein said pharmaceutical composition is useful for the treatment of inflammatory  
5 disease; cancer; central nervous system developmental disease; HIV; FIV; vasculature development disease; haematopoiesis and other chemokine mediated diseases or disorders.

31. A pharmaceutical composition comprising a therapeutically effective amount of a compound of: AMD 3100 ((1,1'-[1,4-phenylenebis(methylene)])bis-1,4,8,11-  
10 tetraazacyclotetradecane), wherein said pharmaceutical composition is useful for the treatment of inflammatory disease; organ transplant rejection; cancer; central nervous system developmental disease; HIV; FIV; vasculature development disease; cardiogenesis developmental disease; haematopoiesis and other chemokine mediated diseases or disorders.

15 32. The pharmaceutical composition of claim 28, 29, 30 or 31, wherein said disease is arthritis.

33. The pharmaceutical composition of claim 28, 29, 30 or 31, wherein said disease is multiple sclerosis.

20

34. The pharmaceutical composition of claim 28, 29, 30 or 31, wherein said cancer is associated with: solid tumors; lymphoma; metastatic tumors; glioblastoma tumors; and other carcinoma tumors.

25 35. The pharmaceutical composition of claim 34, wherein said cancer is; non-small cell lung cancer; lung cancer; breast cancer; prostate cancer; and cancer of other organs.

36. A pharmaceutical composition of claim 28, 29, 30 or 31, wherein said disorder is treated by inhibiting or promoting angiogenesis or by inducing stasis of angiogenesis.

30


37. A method for the treatment of a patient comprising, administering a therapeutically effective dosage of the pharmaceutical composition of claim 28, 29, 30 or 31, to a patient with a disease or disorder that responds to said treatment over a period of time sufficient to effectively treat said disease or disorder.

5

38. A method for the prevention of a disease or disorder in a patient comprising, administering a therapeutically effective dosage of the pharmaceutical composition of claim 28, 29, 30 or 31, to a patient over a period of time sufficient to effectively prevent said disease or disorder.



1/9




AMD3465

FIG. 1




AMD3538

FIG. 2



AMD3500


FIG. 3



AMD3499


FIG. 4

2 / 9




AMD3498

FIG. 5




AMD3497

FIG. 6



AMD3516


FIG. 7



AMD3530


FIG. 8

3/9




AMD3517

FIG. 9




AMD3544

FIG. 10



AMD3543


FIG. 11



AMD3529


FIG. 12

4/9




AMD7049

FIG. 13




AMD7050

FIG. 14




AMD7051

FIG. 15




AMD7059


FIG. 16




AMD7063  
FIG. 17




AMD7060  
FIG. 21




AMD7058  
FIG. 18




AMD7061  
FIG. 22




AMD7032  
FIG. 19



AMD3451  
FIG. 23




AMD7048  
FIG. 20




AMD3454  
FIG. 24

6 / 9




AMD3472

FIG. 25




AMD3526

FIG. 26



AMD3100

FIG. 27



AMD3484

FIG. 28

SUBSTITUTE SHEET (RULE 26)

7/9




FIG. 29






FIG. 30






AMD8630

FIG. 31




AMD7097

FIG. 32




AMD8631

FIG. 33




AMD-Exp 1

FIG. 34




AMD7450

FIG. 35




AMD-Exp 2

FIG. 36



AMD7463

FIG. 37



AMD-Exp 3

FIG. 38

# INTERNATIONAL SEARCH REPORT

Internal Application No

PCT/CA 99/00619

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D257/02 C07D401/12 C07D409/12 C07D403/12 C07D471/06  
C07D401/06 C07D473/00 A61K31/395 //(C07D471/06,257:00,  
221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 747 368 A (JOHNSON MATTHEY PLC) 11 December 1996 (1996-12-11) the whole document & US 5 817 807 A cited in the application ---	1-38
Y	WO 93 12096 A (JOHNSON MATTHEY PLC) 24 June 1993 (1993-06-24) claims & US 5 583 131 A cited in the application ---	1-38
Y	WO 95 18808 A (JOHNSON MATTHEY PLC ;BRIDGER GARY JAMES (US); PADMANABHAN SREENIVA) 13 July 1995 (1995-07-13) claims & US 5 698 546 A cited in the application ---	1-38
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

14 October 1999

Date of mailing of the international search report

02/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 99/00619

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 16494 A (JOHNSON MATTHEY PLC) 1 October 1992 (1992-10-01) claims ---	1-38
A	BRIDGER G J ET AL: "SYNTHESIS AND STRUCTURE-ACTIVITY RELATIONSHIPS OF PHENYLENEBIS(METHYLENE)-LINKED BIS-TETRAAZAMACROCYCLES THAT INHIBIT HIV REPLICATION. EFFECTS OF MACROCYCLIC RING SIZE AND SUBSTITUENTS ON THE AROMATIC LINKER" JOURNAL OF MEDICINAL CHEMISTRY, vol. 38, no. 2, 20 January 1995 (1995-01-20), pages 366-378, XP000196379 ISSN: 0022-2623 the whole document ---	1-38
A	JOAO H C ET AL: "QUANTITATIVE STRUCTURAL ACTIVITY RELATIONSHIP STUDY OF BIS-TETRAAZACYCLIC COMPOUNDS. A NOVEL SERIES OF HIV-1 AND HIV-2 INHIBITORS" JOURNAL OF MEDICINAL CHEMISTRY., vol. 38, no. 19, 1995, pages 3865-3873, XP002118783 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document -----	1-38

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA 99/00619

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 22-25, 27, 37-38  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 22-25, 27, 37-38  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 99/00619

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0747368 A	11-12-1996	AU 5245596 A	19-12-1996
		CA 2178272 A	07-12-1996
		JP 9012560 A	14-01-1997
		NO 962325 A	09-12-1996
		NZ 286711 A	26-02-1998
		US 5817807 A	06-10-1998
		ZA 9604548 A	12-12-1996
WO 9312096 A	24-06-1993	AU 661086 B	13-07-1995
		AU 3165593 A	19-07-1993
		CA 2125978 A	24-06-1993
		CZ 9401188 A	18-10-1995
		EP 0619813 A	19-10-1994
		FI 942849 A	15-06-1994
		HU 67544 A	28-04-1995
		IL 103984 A	30-10-1998
		JP 7501816 T	23-02-1995
		MX 9207316 A	01-06-1993
		NO 942254 A	15-06-1994
		NZ 246179 A	26-03-1996
		PL 173643 B	30-04-1998
		US 5583131 A	10-12-1996
		ZA 9209632 A	18-06-1993
WO 9518808 A	13-07-1995	AU 681478 B	28-08-1997
		AU 1324595 A	01-08-1995
		CA 2179530 A	13-07-1995
		EP 0739345 A	30-10-1996
		FI 962810 A	10-07-1996
		HU 74429 A	30-12-1996
		JP 9509407 T	22-09-1997
		NO 962899 A	10-07-1996
		NZ 277859 A	27-07-1997
		US 5698546 A	16-12-1997
WO 9216494 A	01-10-1992	AU 1372092 A	21-10-1992
		CA 2106068 A	16-09-1992
		FI 933828 A	01-09-1993
		JP 6505728 T	30-06-1994

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**